



New validated *Eucalyptus* SSR markers located in candidate genes involved in growth and plant development

Cintia-Vanesa Acuña^{1*}, Juan-Gabriel Rivas¹, Natalia-Cristina Aguirre¹, Pamela-Victoria Villalba¹, María-Carolina Martínez¹, Martín-Nahuel García¹, Horacio-Esteban Hopp^{1,2} and Susana-Noemí Marcucci-Poltri¹.

¹Instituto de Agrobiotecnología y Biología Molecular (LABiMo), Instituto Nacional de Tecnología Agropecuaria (INTA), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Instituto de Biotecnología, Centro de Investigación en Ciencias Veterinarias y Agronómicas, INTA, N. Repetto y de Los Reseros S/N, Hurlingham B1686IGC, Buenos Aires, Argentina. ²Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, Argentina

Abstract

Aim of study: To validate and characterize new microsatellites or Simple Sequence Repeats (SSR) markers, located within genomic transcribed sequences related to growth and plant developmental traits, in *Eucalyptus* species.

Area of study: *Eucalyptus* species from different Australian origins planted in Argentina.

Material and methods: In total, 134 SSR in 129 candidate genes (CG-SSR) involved in plant development were selected and physically mapped to the *E. grandis* reference genome by bioinformatic tools. Experimental validation and polymorphism analysis were performed on 48 individuals from *E. grandis* and interspecific hybrids (*E. grandis* x *E. camaldulensis*; *E. grandis* x *E. tereticornis*), *E. globulus*, *E. maidenii*, *E. dunnii* and *E. benthamii*.

Main results: 131 out of 134 CG-SSR were mapped on the 11 chromosomes of *E. grandis* reference genome. Most of the 134 analyzed SSR (> 75%) were positively amplified and 39 were polymorphic in at least one species. A search of annotated genes within a 25 kbp up and downstream region of each SSR location retrieved 773 genes of interest.

Research highlights: The new validated and characterized CG-SSR are potentially suitable for comparative QTL mapping, molecular marker-assisted breeding (MAB) and population genetic studies across different species within *Symphyomyrtus* subgenus.

Keywords: CG-SSR; cross-transferability; EST; eucalypts; microsatellite.

Authors' contributions: Conceptualization: HEH, SNMP, CVA. Methodology: CVA, JGR, NCA, PVV. Formal analysis: CVA, JGR, NCA, PVV, MNG. Resources: SNMP, HEH. Writing original draft preparation: CVA. Writing-critical revision of the manuscript: MCM, MNG, NCA, PVV, SNMP, HEH. Project administration and funding acquisition: SNMP, MCM, HEH.

Citation: Acuña, C.V., Rivas, J.G., Aguirre, N.C., Villalba, P.V., Martínez, M.C., García, M.N., Hopp, H.E., Marcucci-Poltri, S.N. (2020). New validated *Eucalyptus* SSR markers located in candidate genes involved in growth and plant development. *Forest Systems*, Volume 29, Issue 3, eSC08. <https://doi.org/10.5424/fs/2020293-17074>.

Supplementary material: Tables S1 and S2 accompany the paper on FS website.

Received: 19 Jun 2020. **Accepted:** 29 Dec 2020.

Copyright © 2020 INIA. This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC-by 4.0) License.

Funding: FONCYT: PICT-2014-0795, PICT 2017-0938-INTA-PNFOR 1104064.

Competing interests: The authors have declared that no competing interests exist.

Correspondence should be addressed to Cintia Vanesa Acuña: acuna.cintia@inta.gob.ar

Introduction

The *Eucalyptus* genus groups forest species and hybrids widely cultivated in the world as forestry plantations for the wood industry as renewable sources for timber, paper and pulp production (Govindan, 2005; Bauhus *et al.*, 2010). The high number of species within this genus, with different agroecological requirements and wood quality characteristics, makes *Eucalyptus* a very valuable resource for its adaptation to the numerous ecosystems worldwide. In this context, the *Eucalyptus* global production is estimated at 20 million hectares (Wingfield *et al.*, 2015).

Over the last 20 years, forest breeding programs, which require long developmental periods, have incorporated different molecular tools (Gudeta, 2018). Molecular markers detect differences between individuals directly from the genome and provide extensive discrete data that can be useful for statistical analyses. These tools are suitable to control the genetic traceability during multiplication processes, evaluate the genetic diversity and make predictions of more reliable breeding values (Cappa *et al.*, 2016; Gudeta, 2018).

SSR or microsatellites, short repetitive DNA sequences distributed throughout the genome showing a high level of polymorphism (first reviewed in plants by Powell

et al., 1996), are the most widely used markers for different applications (Hodel *et al.*, 2016).

The development of microsatellite markers in *Eucalyptus* has evolved over the years in parallel with the increased availability of sequence information. To date, different types of neutral microsatellites are being used in *Eucalyptus* and new SSR markers have been developed within the transcribed regions of the genome. In addition, researchers have designed SSR markers after mining the increasingly larger EST (Expressed Sequence Tags) collections deposited in sequence databases (Kirst *et al.*, 2005; Lehouque *et al.*, 2008; Faria *et al.*, 2010; Acuña *et al.*, 2012 (a, b); Zhou *et al.*, 2014; Grattapaglia *et al.*, 2015).

Several strategies have been explored to find loci controlling traits of interest in woody species. Thus, many genomic studies have reported the analysis of genes and transcription factors expressed during wood formation and xylogenesis in *Eucalyptus* (reviewed by Foucart *et al.*, 2006; Paux *et al.*, 2004; Rengel *et al.*, 2009). Besides, for genes controlling plant growth traits several QTL (Quantitative Trait Loci) approaches have been carried out (revised in Gion *et al.*, 2015; Li *et al.*, 2015; Du *et al.*, 2018; Müller *et al.*, 2019; Kainer *et al.*, 2019).

Therefore, the use of already well-established polymorphic markers located in CG for plant traits is an interesting approach for mapping purposes and population genetic studies with an emphasis in non-model species (Acuña *et al.*, 2012b, 2014; Pomponio *et al.*, 2015; Azpilicueta *et al.*, 2016). Also, the availability of *E. grandis* genome sequence, with annotated and classified genes, is an important information source to study and characterize different traits of interest (Myburg *et al.*, 2014).

Although high-throughput sequence-based SNP marker assays are increasingly becoming available (Silva-Junior *et al.*, 2015; Aguirre *et al.*, 2019), microsatellites still constitute a very useful and accessible tool for fast and precise genetic analysis in *Eucalyptus* (Grattapaglia *et al.*, 2015). Besides genetic diversity studies, SSR have numerous uses, including cultivar or clone fingerprinting, population structure, marker-assisted selection, linkage map development and QTL mapping, among others, thus showing an important role in this genomic age (Hodel *et al.*, 2016).

In this study, we in silico characterized and in vitro validated new microsatellite markers located in CG (structural genes and transcriptional factors) related to plant growth and development. The selection of SSR located on these genes was based on data from a previous study of our group (Acuña *et al.*, 2012a), with a focus on genome regions potentially involved in these important characteristics for tree breeding. The identified SSR markers were wet-lab validated in five different *Eucalyptus* species and hybrids, and physically mapped on the *E. grandis* reference genome. Furthermo-

re, we also identified and analyzed known predicted genes contiguous (<25kbp) to these SSR markers.

Materials and Methods

Leaves from 48 individuals of *Eucalyptus* sp. were analyzed: 12 individuals of *Eucalyptus grandis* (4 individuals from a clonal population, 4 individuals from two controlled crosses and 4 individuals from their offspring), *E. globulus* (6 individuals), *E. maidenii* (8 individuals), *E. dunnii* (8 individuals), *E. benthamii* (8 individuals) and the hybrids *E. grandis* x *E. camaldulensis* (3 individuals) and *E. grandis* x *E. tereticornis* (3 individuals). Trees were planted in EEA INTA Concordia (31°22'28.9"S 58°07'01.0"W) and IRB-CIRN-CNIA-INTA (34°36'58.6"S 58°40'06.2"W), Argentina. Total DNA was extracted from young leaves using the CTAB method with modifications, as described in Marcucci Poltri *et al.* (2003).

We selected SSR markers on CG from a previous study (Acuña *et al.*, 2012a), 1,140 SSR within 952 CG were in silico characterized. These CG had been selected for their possible biological function predicted according to Gene Ontology (GO) (Ashburner *et al.*, 2000; <http://www.geneontology.org/>) using Blast2GO (Conesa *et al.*, 2005). From those genes, in the present study, 129 CG with 134 SSR sequences were selected based on their correspondence to genes and transcriptional factors involved in different plant growth and developmental features.

Validation was carried out using PCR reactions in a final volume of 12 µl with 20 ng of genomic DNA, 0.25 µM of each primer (Alpha DNA, Canada), 2mM MgCl₂, 0.2 mM of each dNTP, 1X reaction buffer and 1U Platinum Taq polymerase (Invitrogen, Waltham, USA). Amplifications were performed following a denaturation step of 5 min at 94 °C, 35 cycles of 1 min at 94 °C, 1 min at annealing temperature and 1 min at 72 °C. The final extension step was for 10 min at 72 °C. The SSR amplification products were denatured for 5 min in denaturing loading buffer at 95 °C and separated by a 6% polyacrylamide gel electrophoresis (6% acrylamide/bisacrylamide 20:1, 7.5 M urea, 0.5 × TBE) along with a 25 bp DNA ladder standard (Invitrogen, Waltham, USA). The DNA silver-staining procedure of Promega (Madison, WI, USA) was used for visualization. Details on primer sequences, SSR location, annealing temperature and product sizes are described in Table S1 [supplementary].

We carried out the in silico characterization of the CG-SSR through physical mapping and nearby gene search. The 134 SSR obtained sequences were mapped to the *E. grandis* reference genome (Myburg *et al.*, 2014) (<http://phytozome.jgi.doe.gov>, version 2.0). Mapping was performed using the Bowtie2 alignment tool with default settings (Langmead & Salzberg, 2012). A custom Perl

script was used to determine the annotated genes of the *E. grandis* genome within a flanking region of 50 kbp (up to 25 kbp from each SSR locus). The predicted genes classified and reported by Myburg *et al.* (2014) were used to describe some of the genes found within the window surrounding each SSR.

Results and Discussion

Selected CG-SSR markers

Based on the results from a previous study (Acuña *et al.*, 2012a), we selected sequences similar to structural genes and transcriptional factors involved in plant developmental features and obtained 129 CG with 134 SSR sequences (details of SSR markers in Table S1 [suppl.]). This study revealed the following distribution of the selected 129 GC within the GO terms: most of them belonged to the “Biological Process” category and within this category, to the subcategories “Metabolic Process of Organic Substances” (17%), “Cellular Metabolic Process” (15%) and “Primary Metabolic Process” (15%), among other less represented subcategories. Also, most of the GO terms within the “Molecular Function” class belonged to the subcategories “Binding to Heterocyclic Compounds” (15%), “Binding to Organic Cyclic Compounds” (15%) and “Ion Binding” (14%). Among them, we detected SSR in transcriptional factor genes involved in xylogenesis (MYB, bZIP, WRKY, SWI/SNF, ARF) (Rengel *et al.*, 2009) and responses to abiotic stress (BES, bZIP) (Bechtold & Field, 2018) (Table S1 [suppl.]).

Marker validation in different *Eucalyptus* spp.

Most of the SSR (75.4%) analyzed in the laboratory were positive PCR-amplified according to similar studies in *Eucalyptus* (Acuña *et al.*, 2012a; He *et al.*, 2012; Zhou *et al.*, 2014). Nonspecific amplicons (17 SSR) or amplicons with sizes above 500 bp (20 SSR) were discarded. Thus, 64 markers resulted in amplification products of the expected size according to bioinformatic analysis. Among them, 39 (29%) were polymorphic (*i.e.*, they had at least 2 alleles in at least one species) and 25 were monomorphic (Table S1 [suppl.]). The polymorphism rate of EST-SSR was similar to that described by Faria *et al.* (2011) (25%) and Acuña *et al.* (2012a) (30%), but lower than that observed by Faria *et al.* (2010) (39%) and Grattapaglia *et al.* (2015) (65%) in *Eucalyptus*.

The relative proportions of repeated motifs in polymorphic SSR were 28.2% for di-, 56.4% for tri-, 10.2% for tetra-, and 5.2% for pentanucleotides. Our results are similar to those reported by other authors, in which trinucleotide repeats were the most common, followed by

di- and tetranucleotide repeats (Varshney *et al.*, 2005, Grattapaglia *et al.*, 2015).

The number of alleles per marker (between 2 and 7) (Table S1 [suppl.]) is equivalent to that reported by other authors for EST-SSR markers validated on a small number of samples (about 8 individuals per species) (Zhou *et al.*, 2014). On the other hand, the values found in the present study are lower than those described by Faria *et al.* (2010, 2011) and Grattapaglia *et al.* (2015) in *Eucalyptus*. These results could be explained by the marker selection criteria used. While these studies based their selection on the polymorphism level, we selected SSR markers focusing on their putative function in growth and plant development. Nevertheless, the number of alleles per marker may increase with a larger sample size.

Physical mapping and nearby gene search

The alignment of the 134 CG-SSR sequences against the *E. grandis* public reference genome revealed that 131 SSR were mapped on the 11 chromosomes, while three of the markers were located on scaffolds. The number of SSR by chromosome ranged between 4 (Chromosome 5) and 17 (chromosomes 3, 8 and 11), thus showing a good distribution in the genome (Table S2 [suppl.]).

According to an exhaustive bibliographic revision of the available publications that developed this kind of markers in *Eucalyptus*, only 16 of the 134 SSR validated here coincide with those of other studies (2 in Yasodha *et al.*, 2008; 8 in Rengel *et al.*, 2009; 4 in He *et al.*, 2012; 4 in Zhou *et al.*, 2014 and 1 in Grattapaglia *et al.*, 2015, where some markers were shared between studies). Nonetheless, none of these studies involved the characterization related to plant growth and developmental traits. Moreover, only in this study and in that by Grattapaglia *et al.* (2015), EST-SSR were aligned to the *E. grandis* genome sequence, thus providing information on their distribution and physical position (Table S1 [suppl.]).

Interestingly, the 39 polymorphic SSR markers are located in protein-coding CG, *e.g.* serine-threonine kinase (which is involved in the completion of embryonic development in dormant seeds), F-box type (signal transduction and cell cycle) (Jia *et al.*, 2020) and various transcription factors that regulate processes of cellular development, seed maturation, floral development, among others (bZIP, GATA, BES1) (Bechtold and Field, 2018) (Table S1 [suppl.]).

Additionally, we performed a search for genes of interest that could be linked to the identified SSR within a flanking window of 50 kbp (25kbp up- and downstream regions). This window size was selected based on Linkage Disequilibrium (LD) in *E. grandis* reported by Silva-Junior *et al.*, 2015). This analysis yielded 773 *E. grandis* predicted genes (named Eucgr. in Myburg *et al.*, 2014)

neighbouring these SSR. Among them, 394 were within a Gene Ontology (GO) category (Table S2 [suppl.]).

Based on Myburg *et al.* (2014), who classified predicted genes according to different classes related to wood quality, 30 of the 773 genes belong to the following categories: 3 into “MYB Transcription Factors”, 1 into “Genes Encoding Laccases and Peroxidases”, 8 into “Lignin Biosynthesis”, 7 into “predicted cellulose and xylan genes” and 11 into “Interpro Domain of 968 Unique *Eucalyptus* Genes”. This categorization gives these markers an added value, since we detected genes related not only to plant growth and development, but also to wood quality (Table S2 [suppl.]). Examples of genes related to wood quality are cinnamoyl CoA reductase (CCR), phenylalanine ammonia-lyase (PAL), 4-coumarate-CoA ligase (4CL) and Caffeic Acid O-Methyl Transferase (COMT) genes, which code for the key enzymes in lignin biosynthesis (Boerjan *et al.*, 2003). Other of these identified genes are PARVUS, cellulose synthase (CESA) and sucrose synthase (SUSY), which are involved in cellulose and xylan biosynthesis (Myburg *et al.*, 2014).

In this work, the evaluated candidate genes sequences were up to 25 kbp distance from the validated SSR markers. Therefore, linkage between them seems to be high enough to make this panel of SSR markers useful in future association mapping studies for *Eucalyptus* breeding purposes.

Conclusions

In the present study, a new set of SSR especially located in candidate genes for growth and plant development is proposed as a tool for *Eucalyptus* genetic analysis. Additionally, some of the SSR are particularly interesting, because they are close to candidate genes for wood quality.

These new CG-SSR markers, in addition to those already publicly available, could be included in studies for the identification of different *Eucalyptus* genetic materials, in population genetics, taxonomy, verification of synteny and collinearity between different *Eucalyptus* maps. Furthermore, they could be implemented in QTL and association mapping studies and genomic selection through relatedness correction in breeding value predictions.

Acknowledgements

The authors would like to thank Pablo Pathauer, Javier Oberschelp, Leonel Harand, Mauro Surenciski and Martín Marcó for providing the plant material. Also, we like to express our gratitude to Janet Higgins who developed the Perl script used on this paper and a sincere thank you

to Julia Sabio y García for her diligent proofreading of the manuscript.

References

- Acuña C, Fernandez P, Villalba P, García M, Hopp E, Marcucci Poltri S, 2012a. Discovery, validation and in silico functional characterization of EST-SSR markers in *Eucalyptus globulus*. *Tree Genet Genomes* 8:289-301. <https://doi.org/10.1007/s11295-011-0440-0>
- Acuña C, Villalba P, Pathauer P, Hopp E, Marcucci Poltri S, 2012b. Characterization of novel microsatellite markers in candidate genes for wood properties for application in functional diversity assessment in *Eucalyptus globulus*. *Electron J Biotechnol* 15 (2): 12-28. <https://doi.org/10.2225/vol15-issue2-fulltext-3>
- Acuña C, Villalba P, Hopp E, Marcucci Poltri S, 2014. Transferability of microsatellite markers located in candidate genes for wood properties between *Eucalyptus* species. *Forest Systems* 2014 23(3): 506-512. <https://doi.org/10.5424/fs/2014233-05279>
- Aguirre N, Filippi C, Zaina G, Rivas J, Acuña C, Villalba P, García M, González S, Rivarola M, Martínez M, *et al.*, 2019. Optimizing ddRADseq in Non-Model Species: A Case Study in *Eucalyptus dumii* Maiden. *Agronomy* 2019, 9, 484. <https://doi.org/10.3390/agronomy9090484>
- Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, *et al.*, 2000. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet* 25:25-29. <https://doi.org/10.1038/75556>
- Azpilicueta MM, El Mujtar VA, Gallo L, 2016. A Searching for molecular insight on hybridization in *Nothofagus* spp. forests at Lagunas de Epulauquen, Argentina. *Bosque* 2016, 37(3), 591-601. <https://doi.org/10.4067/S0717-92002016000300016>
- Bauhus J, van der Meer P and Kanninen M, 2010. Ecosystem goods and services from plantation forests. London, Great Britain: Earthscan. 254 pp. (Earthscan forest library). <https://doi.org/10.4324/9781849776417>
- Bechtold U, Feld B, 2018. Molecular mechanisms controlling plant growth during abiotic stress. *J Exp Bot* 69 (11): 2753-2758. <https://doi.org/10.1093/jxb/ery157>
- Boerjan W, Ralph J, Baucher M, 2003. Lignin Biosynthesis. *Annu Rev Plant Biol* 54 (1):519-546. <https://doi.org/10.1146/annurev.arplant.54.031902.134938>
- Cappa EP, Klápště J, Garcia, MN, Villalba PV, Marcucci Poltri, SN, 2016. SSRs, SNPs and DArTs comparison on estimation of relatedness and genetic parameters: precision from a small half-sib sample population of *Eucalyptus grandis*. *Mol Breeding* 36:97. <https://doi.org/10.1007/s11032-016-0522-7>

- Conesa A, Götz S, García-Gómez JM, Terol J, Talón M, Robles M, 2005. Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* 21:3674-3676. <https://doi.org/10.1093/bioinformatics/bti610>
- Du Q, Lu W, Quan M, Xiao L, Song F, Li P, Zhou D, Xie J, Wang L, Zhang D, 2018. Genome-Wide Association Studies to Improve Wood Properties: Challenges and Prospects. *Front Plant Sci* 29: 1912. <https://doi.org/10.3389/fpls.2018.01912>
- Faria DA, Mamani EMC, Pappas MR, Pappas jr GJ, Grattapaglia D, 2010. A selected set of EST-derived microsatellites, polymorphic and transferable across 6 species of *Eucalyptus*. *J Hered* 101: 512-520. <https://doi.org/10.1093/jhered/esq024>
- Faria DA, Mamani EMC, Pappas GJ, Grattapaglia D, 2011. Genotyping systems for *Eucalyptus* based on tetra-, penta-, and hexanucleotide repeat EST microsatellites and their use for individual fingerprinting and assignment tests. *Tree Genet Genomes* 7, 63-77. <https://doi.org/10.1007/s11295-010-0315-9>
- Foucart C, Paux E, Ladouce N, San-Clemente H, Grima-Pettenati J, Sivadon P, 2006. Transcript profiling of a xylem vs phloem cDNA subtractive library identifies new genes expressed during xylogenesis in *Eucalyptus*. *New Phytol* 170: 739-752. <https://doi.org/10.1111/j.1469-8137.2006.01705.x>
- Gion J, Chaumeil P, Plomion C, 2015. EucaMaps: linking genetic maps and associated QTLs to the *Eucalyptus grandis* genome. *Tree Genet Genomes* 11, 795. <https://doi.org/10.1007/s11295-014-0795-0>
- Govindan M, 2005. *Eucalyptus: the Genus Eucalyptus*. Edited by John J. W. Coppen (Natural Resources Institute, University of Greenwich, UK). Taylor and Francis, London. 2002. ISBN 0-415-27879-1. *J. Nat. Prod.* 68: 151-152. <https://doi.org/10.1021/np0307789>
- Grattapaglia D, Mamani E, Silva-Junior O, Faria D, 2015. A novel genome-wide microsatellite resource for species of *Eucalyptus* with linkage-to-physical correspondence on the reference genome sequence. *Mol Ecol Resour* 15 (2): 437-448. <https://doi.org/10.1111/1755-0998.12317>
- Gudeta TB, 2018. Molecular marker based genetic diversity in forest tree populations. *Forest Res Eng Int J.* 18;2(4):176-182. <https://doi.org/10.15406/freij.2018.02.00044>
- He X, Wang Y, Li F, Weng Q, Li M, Xu L A, Gan S, 2012. Development of 198 novel EST-derived microsatellites in *Eucalyptus* (Myrtaceae). *Am J Bot* 99(4), e134-e148. <https://doi.org/10.3732/ajb.1100442>
- Hodel RGJ, Segovia-Salcedo MC, Landis JB, Crowl AA, Sun M, Liu X, Gitzendanner MA, Douglas NA, Germain-Aubrey CC, Chen S, Soltis, D E, Soltis PS, 2016. The report of my death was an exaggeration: A review for researchers using microsatellites in the 21st century. *Appl Plant Sci* 4(6): 1600025. <https://doi.org/10.3732/apps.1600025>
- Jia Z, Zhao B, Liu S, Lu Z, Chang B, Jiang H, Cui H, He Q, Li W, Jin B, Wang L, 2020. Embryo transcriptome and miRNA analyses reveal the regulatory network of seed dormancy in *Ginkgo biloba*. *Tree Physiol.* tpa023. <https://doi.org/10.1093/treephys/tpaa023>
- Kainer D, Padovan A, Degenhardt J, Krause S, Mondal P, Foley WJ, Külheim C, 2019. High marker density GWAS provides novel insights into the genomic architecture of terpene oil yield in *Eucalyptus*. *New Phytol.* 223: 1489-1504. <https://doi.org/10.1111/nph.15887>
- Kirst M, Cordeiro CM, Rezende G, Grattapaglia D, 2005. Power of microsatellite markers for fingerprinting and parentage analysis in *Eucalyptus grandis* breeding populations. *J Hered* 96(2):161-166. <https://doi.org/10.1093/jhered/esi023>
- Langmead B, Salzberg SL, 2012. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* 9: 357. <https://doi.org/10.1038/nmeth.1923>
- Lehouque G, Sanhueza R, Melo F, 2008. Development of MultiTAAG: an Automated Genotyping System for *Eucalyptus* Species Using Multiplex Amplification of Microsatellite Markers. *Boletín del CIDEU* 6-7: 25-34.
- Li F, Zhou C, Weng Q, Li M, Yu X, Guo Y, Wang Y, Zhang X, Gan, S, 2015. Comparative genomics analyses reveal extensive chromosome colinearity and novel quantitative trait loci in *Eucalyptus*. *PloS one*, 10(12), e0145144. <https://doi.org/10.1371/journal.pone.0145144>
- Marcucci Poltri SN, Zelener N, Rodriguez Traverso J, Gelid P, Hopp HE, 2003. Selection of a seed orchard of *Eucalyptus dumonii* based on genetic diversity criteria calculated using molecular markers. *Tree Physiol* 23(9): 625-632. <https://doi.org/10.1093/treephys/23.9.625>
- Müller BSF, de Almeida Filho JE, Lima BM, García CC, Missiaggia A, Aguiar AM, Takahashi E, Kirst M, Gezan SA, Silva-Junior OB, *et al.*, 2019. Independent and Joint GWAS for growth traits in *Eucalyptus* by assembling genome: wide data for 3373 individuals across four breeding populations. *New Phytol*, 221: 818-833. <https://doi.org/10.1111/nph.15449>
- Myburg A, Grattapaglia D, Tuskan G, Hellsten U, Hayes RD, Grimwood J, Jenkins J, Lindquist E, Tice H, Bauer D, *et al.*, 2014. The genome of *Eucalyptus grandis*. *Nat* 510: 356-362.
- Paux E, Tamasloukht M, Ladouce N, Sivadon P, Grima-Pettenati J, 2004. Identification of genes preferentially expressed during wood formation in *Eucalyptus*. *Plant Mol Biol* 55: 263-80. <https://doi.org/10.1007/s11103-004-0621-4>
- Pomponio M, Acuña C, Petreath VL, Lauenstein D, Marcucci Poltri S, Torales S, 2015. Characterization

- of functional SSR markers in *Prosopis alba* and their transferability across *Prosopis* species. *Forest Systems*, 24(2), eRC04. <https://doi.org/10.5424/fs/2015242-07188>
- Powell W, Machray GC, Provan J, 1996. Polymorphisms revealed by simple sequence repeats. *Trends Plant Sci* 1 (7): 215-222. [https://doi.org/10.1016/S1360-1385\(96\)86898-0](https://doi.org/10.1016/S1360-1385(96)86898-0)
- Rengel D, Clemente HS, Servant F, Ladouce N, Paux E, Wincker P, Couloux A, Sivadon P, Grima-Pettenati J, 2009. A new genomic resource dedicated to wood formation in *Eucalyptus*. *BMC Plant Biol* 9: 36. <https://doi.org/10.1186/1471-2229-9-36>
- Silva-Junior OB, Grattapaglia D, 2015. Genome-wide patterns of recombination, linkage disequilibrium and nucleotide diversity from pooled resequencing and single nucleotide polymorphism genotyping unlock the evolutionary history of *Eucalyptus grandis*. *New Phytol* 208: 830-845. <https://doi.org/10.1111/nph.13505>
- Silva-Junior OB, Faria DA, Grattapaglia D, 2015. A flexible multi-species genome-wide 60K SNP chip developed from pooled resequencing of 240 *Eucalyptus* tree genomes across 12 species. *New Phytol.* 2015, 206, 1527-1540. <https://doi.org/10.1111/nph.13322>
- Varshney RK, Graner A, Sorrells ME, 2005. Genic microsatellite markers in plants: features and applications. *Trends Biotechnol.* 23: 48-55. <https://doi.org/10.1016/j.tibtech.2004.11.005>
- Wingfield MJ, Brockerhoff EG, Wingfield BD, Slippers B, 2015. Planted forest health: The need for a global strategy. *Sci* 349: 832-836. <https://doi.org/10.1126/science.aac6674>
- Yasodha R, Sumathi R, Chezhian P, Kavitha S, Ghosh M, 2008. *Eucalyptus* microsatellites mined in silico: survey and evaluation. *J Genet* 87:21-25. <https://doi.org/10.1007/s12041-008-0003-9>
- Zhou C, He X, Li F, Weng Q, Yu X, Wang Y, Li M, Shi J, Gan S, 2014. Development of 240 novel EST-SSRs in *Eucalyptus* L'Hérit. *Mol Breeding* 33: 221-225. <https://doi.org/10.1007/s11032-013-9923-z>