



Proceeding Paper

Could the Interaction between Arbuscular Mycorrhizal Fungi and Biostimulants Improve the Plant Physiological Status of *Prosopis alba* Seedlings? †

Ana Carolina Santacruz-García ^{1,*} , Adriana Teresita Gomez ², María Gracia Senilliani ³, Analia del Valle Guzmán ³, Mónica Beatriz Sagadin ⁴, Mauricio Ewens ⁵, Rodrigo Sebastian Frías ³, Cristian Coria ³, Joaquín Gómez ³ and Mónica Azucena Nazareno ¹

- ¹ Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Instituto de Ciencias Químicas, Facultad de Agronomía y Agroindustrias, Universidad Nacional de Santiago del Estero, Santiago del Estero C.P. 4200, Argentina
 - ² Estación Experimental Agropecuaria Santiago del Estero, Instituto Nacional de Tecnología Agropecuaria, Santiago del Estero C.P. 4200, Argentina
 - ³ Facultad de Ciencias Forestales, Instituto de Silvicultura y Manejo de Bosques (INSIMA), Universidad Nacional de Santiago del Estero, Santiago del Estero C.P. 4200, Argentina
 - ⁴ Instituto de Fisiología y Recursos Genéticos Vegetales (IFRGV), Centro de Investigaciones Agropecuarias (CIAP), Instituto Nacional de Tecnología Agropecuaria (INTA), Unidad de Estudios Agropecuarios (UDEA) INTA-CONICET, Córdoba C.P. 5119, Argentina
 - ⁵ Estación Experimental Fernández-UCSE (Convenio Universidad Católica Sgo del Estero-Provincia Sgo del Estero), Departamento Robles, Santiago del Estero C.P. 4322, Argentina
- * Correspondence: anacaro.santacruz@gmail.com; Tel.: +54-9-385-474-4264
- † Presented at the 3rd International Electronic Conference on Forests—Exploring New Discoveries and New Directions in Forests, 15–31 October 2022; Available online: <https://iecf2022.sciforum.net/>.



Citation: Santacruz-García, A.C.; Gomez, A.T.; Senilliani, M.G.; Guzmán, A.d.V.; Sagadin, M.B.; Ewens, M.; Frías, R.S.; Coria, C.; Gómez, J.; Nazareno, M.A. Could the Interaction between Arbuscular Mycorrhizal Fungi and Biostimulants Improve the Plant Physiological Status of *Prosopis alba* Seedlings? *Environ. Sci. Proc.* **2022**, *22*, 53. <https://doi.org/10.3390/IECF2022-13089>

Academic Editor: Rodolfo Picchio

Published: 21 October 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: This study aimed to evaluate the synergy between arbuscular mycorrhizal fungi (AMF) and foliar biostimulant applications (phytoextracts) on an important Argentinian forest native species (*Prosopis alba*) during the nursery stage. We tested biochemical parameters (MDA, malondialdehyde, an oxidative stress biomarker, and photosynthetic pigments) on *P. alba* seedlings sprayed with three different phytoextracts and inoculated with mycorrhizal strains of different local origins. Considering that the statistical model was not significant at the preliminary level, we did not observe synergism between the different forms of bioinsumes evaluated by analysing biochemical characteristics. However, regardless of AMF inoculation, plants with foliar applications of *Larrea divaricata* at 3% w/v showed a lower accumulation of the oxidative stress biomarker, MDA and a lower total carotenoid content ($p > 0.1$). Although there were no significant differences, trends indicate positive relationships between neck diameter and mycorrhizal response in plants treated with M1 with foliar applications of *Larrea divaricata* at 3% w/v. However, height does not consistently respond to mycorrhizae interacting with biostimulants. Consequently, deeper analysis is needed to understand the effect of the interaction between AMF with biostimulants for improving the plant's physiological status. Deepening research in this regard will result in significant benefits for restoration activities.

Keywords: algarrobo blanco; Chaco; mycorrhizae; plant physiological status; AMF

1. Introduction

Mycorrhizae and biostimulants are among the most innovative biological techniques used in large-scale food production. Positive effects on plant growth have been reported. However, the use of these “biofertilizers” in afforestation is limited. Currently, there is growing market interest in bioinsumes for agribusiness. This is due to the need to promote sustainable production techniques that efficiently use their resources [1,2]. Bioinsumes are

natural extracts or microorganisms that could act on plant physiology, improving their tolerance to stress, nutritional efficiency, and/or quality characteristics [1,3]. An example of these products is *biocontrol* agents. These products are mainly derived from beneficial fungi, which, through symbiotic relationships, protect plants against pathogen attack [4,5]. Another class of bioinsumes are *biostimulants*. These products promote germination, growth, flowering and/or fruit development, despite not being nutrients, soil improvers or pesticides. These kinds of products are characterised by improving tolerance to plant abiotic stress [1,6]. Natural extracts have great potential as biostimulants; however, their activity is still little known [1]. Interestingly, numerous studies highlight the protective effect of bio-inputs against stress factors such as drought, salinity, and pathogen attack, among others [3,7,8]. Deepening research about the interaction between different bioinsumes could represent an effective strategy to enhance plant responses to oxidative stress.

2. Methodology

2.1. Plant Material and Experimental Conditions

We conducted an experiment from November 2021 to February 2022 in the Experimental Station “Fernández” (Agreement Catholic University of Santiago del Estero-Province of Santiago del Estero) in Santiago del Estero, Argentina ($-27^{\circ}56'0''$ S, $65^{\circ}52'50''$ W). Seedlings were produced in trays of individual cells in a nursery with 50% of shading under natural light conditions for 45 days. After that, the plants were exposed to full sun in the acclimation phase until completing 90 days.

2.2. Isolation, Multiplication and Application of AMF Inocula, and Preparation and Application of Biostimulants

We selected mixed inocula of native AMF from *P. alba* stands located in the Argentine Chaco Region, with two different locations: Padre Lozano (M1) in the Western Chaco Domain, and Colonia Benítez (M2) in the Eastern Chaco Domain, Chaco province, according to the methodology proposed by Sagadin et al. (2018) [9]. M1: *Claroideoglossum claroideum*, *Claroideoglossum etunicatum*; *Diversispora spurca*, *Funneliformis mosseae*, and *Rhizophagus intraradices*; M2: *Claroideoglossum claroideum*, *Claroideoglossum etunicatum*, *Funneliformis constrictum*, *Funneliformis mosseae*, and *Rhizophagus clarus*. SM: Non-inoculated. The inoculation was performed at sowing by applying 10 g of AMF inoculum per container for each inoculum in the planting hole Salto et al. (2020) [10].

Biostimulants were prepared and applied according to the methodology described in Santacruz-García et al. (2022) [11]. J1Y1 biostimulant is a mixture of both species (*Larrea divaricata* and *Ilex paraguariensis*, 1% w/v: 1% w/v). Foliar applications were made twice during the acclimation stage (on days 7 and 14). The measurements were made on the 21st day of the acclimation stage.

2.3. Plant Biochemical Responses

In this study, (i) malondialdehyde (MDA), an oxidative stress biomarker, and (ii) photosynthetic pigments, total contents of chlorophylls and carotenoids, were evaluated according to methodology followed by Santacruz-García et al. (2022) [11]. Leaf samples were collected by triplicate, evaluating three plants per treatment.

2.4. Plant Morphological Responses

For this assay, we considered the following morphological characteristics: (i) stem neck diameter (SND) and (ii) shoot height (SH), according to standard methodologies proposed by Santacruz-García et al. (2022) [11]. Measurements were made on ten plants per treatment.

2.5. Mycorrhizal Response

Mycorrhizal Response (MR) was calculated for (i) stem neck diameter (SND) and (ii) shoot height (SH) according to the relationship described by Cavagnaro et al. (2003) [12]:

$$MR = \frac{M - \text{mean SM}}{\text{mean SM}} 100$$

M corresponds to the morphological characteristic evaluated and mean SM corresponds to the non-inoculated plants.

2.6. Statistical Analysis

For assessments of biochemical responses to the biostimulant application, data were analysed through a mixed linear model (MM), using factors (biostimulant, AMF inoculum, and their interactions) as fixed effects. The individual plant was considered a random effect. For the analysis of the morphological variables (SND and SH), a mixed linear model was used using as a random effect the repetition by plot (AMF inoculum_Bioestimulant). While the mycorrhizal response (MR) of SND and SH was analysed with a mixed general model using heteroscedasticity of variances in repetition per plot. The statistical software used was Infostat/2017 (InfoStat Group V.2017, Cordoba, Argentina) with an $\alpha = 0.05$.

3. Results and Discussion

3.1. Plant Biochemical Responses

We did not observe synergism between the different forms of bioinsumes evaluated considering biochemical characteristics. According to the statistical analysis, the interactions between treatments significantly did not affect the plant’s biochemical responses (Table 1). These results could be explained considering that the colonization efficiency of AMF is highly related to the environmental conditions and plant genotypes [13]. It is probable that the interaction between the selected biostimulants and the AMF strains used in this study did not enhance plant growth and nutrition. It is necessary to explore different AMF species to evaluate their synergist potential with biostimulants to enhance the production of *P. alba* seedlings [14].

Table 1. Mean and standard deviation for the biochemical variables: total contents of chlorophylls ($\mu\text{g g}^{-1}$ FW), carotenoids ($\mu\text{g g}^{-1}$ FW), and malondialdehyde (MDA, nmol g^{-1} FW). Treatments are combination of inoculation (SM: Non inoculated, M1: *Claroideoglo mus claroideum*, *Claroideoglo mus etunicatum*; *Diversispora spurca*, *Funneliformis mosseae*, and *Rhizophagus intraradices*; M2: *Claroideoglo mus claroideum*, *Claroideoglo mus etunicatum*, *Funneliformis constrictum*, *Funneliformis mosseae*, and *Rhizophagus clarus*) and biostimulant foliar applications (SB: Seedlings sprayed with only water, J3: seedlings sprayed with *Larrea divaricata* (3% w/v), Y2: seedlings sprayed with *Ilex paraguariensis* (2% w/v), and J1Y1: mixture of both biostimulants (1% w/v: 1% w/v). Stars indicate the significance level. Significance levels: *** <0.001; ** <0.01; * <0.05; <0.1.

Treatment	Chlorophylls	Carotenoids	MDA
SMSB	2151.90 ± 160.16	372.13 ± 12.46	28.84 ± 1.53
SMJ3	1880.24 ± 129.30	326.79 ± 29.06	26.20 ± 0.99
SMY2	2014.71 ± 63.37	349.14 ± 4.64	29.07 ± 3.30
SMJ1Y1	2140.50 ± 139.36	373.87 ± 26.33	29.18 ± 3.50
M1SB	2161.63 ± 122.16	366.99 ± 20.26	28.10 ± 2.79
M1J3	2018.92 ± 179.92	344.04 ± 6.95	26.85 ± 3.63
M1Y2	2070.84 ± 368.22	361.34 ± 59.37	28.65 ± 4.83
M1J1Y1	2375.28 ± 195.11	396.44 ± 80.80	29.41 ± 1.15
M2SB	2070.59 ± 104.51	366.92 ± 1.75	30.83 ± 2.37
M2J3	2078.85 ± 555.13	348.94 ± 97.71	25.09 ± 2.63
M2Y2	2125.39 ± 253.52	379.42 ± 33.70	26.96 ± 1.21
M2J1Y1	2375.28 ± 195.11	412.21 ± 25.35	27.99 ± 5.85

However, foliar applications of biostimulants showed a slightly significant effect on the total content of carotenoids and MDA ($p < 0.1$, Table 2). Seedlings sprayed with *Larrea divaricate* (3% w/v) exhibited lower values in both biochemical characteristics, which could be related to the antioxidant effect of this biostimulant. These results could be associated

with the preventive action of this biostimulant application against the oxidative stress that affects different physiological processes [7]. It is interesting to highlight that our results confirmed the observed effects of *L. divaricata* (3% w/v) as a potential biostimulant of *P. alba* during the acclimation stage [11].

Table 2. Mean and standard deviation for the biochemical variables: total contents of chlorophylls ($\mu\text{g g}^{-1}$ FW), carotenoids ($\mu\text{g g}^{-1}$ FW), and malondialdehyde (MDA, nmol g^{-1} FW). Treatments are biostimulant foliar applications (SB: Seedlings sprayed with only water, J3: seedlings sprayed with *Larrea divaricata* (3% w/v), Y2: seedlings sprayed with *Ilex paraguariensis* (2% w/v), and J1Y1: mixture of both biostimulants (1% w/v: 1% w/v). Different letters indicate significant differences, according to LSD Fisher pairwise comparison procedure with α : 0.1. Stars indicate the significance level. Significance levels: **** <0.001; *** <0.01; ** <0.05; * < 0.1.

Treatment	Chlorophylls	Carotenoids *	MDA *
SB	2128.04 ± 121.44 a	369.68 ± 12.14 ab	29.26 ± 2.33 b
J3	1992.67 ± 311.59 a	339.92 ± 52.07 a	26.09 ± 2.42 a
Y2	2070.31 ± 230.72 a	363.30 ± 36.67 ab	28.23 ± 3.14 ab
J1Y1	2244.57 ± 297.77 a	394.17 ± 47.38 b	28.86 ± 3.52 ab

3.2. Plant Morphological Responses

The diameter and height of the seedlings did not show significant differences with the application of mycorrhizae and biostimulants in interaction, nor with the fixed effects. However, there are trends in favour of M1 and J3. However, there are positive trends between the SND with the application of J3 and M1, while for the variable height is not so evident this trend in the interaction between AMF inoculum and biostimulant applications. (Table 3). This may be related to the fact that the diameter is a conduction tissue and the inoculum M1 comes from an arid site could favour this seedling feature [15].

Table 3. Mean and standard error of Stem neck diameter (SND) and Stem height (SH) of *Prosopis alba* seedlings Interaction are combination of inoculation (SM: Non inoculated, M1: *Claroideoglopus claroideum*, *Claroideoglopus etunicatum*; *Diversispora spurca*, *Funneliformis mosseae*, and *Rhizophagus intraradices*; M2: *Claroideoglopus claroideum*, *Claroideoglopus etunicatum*, *Funneliformis constrictum*, *Funneliformis mosseae*, and *Rhizophagus clarus*) and biostimulant foliar applications (SB: Seedlings sprayed with only water, J3: seedlings sprayed with *Larrea divaricata* (3% w/v), Y2: seedlings sprayed with *Ilex paraguariensis* (2% w/v), and J1Y1: mixture of both biostimulants (1% w/v: 1% w/v).

Treatment	SND (mm)	SH (cm)
M1 J3	3.60 ± 0.1	38.62 ± 1.39
M1 SB	3.54 ± 0.13	38.69 ± 1.37
SM SB	3.53 ± 0.07	36.15 ± 1.34
SM J1Y1	3.52 ± 0.07	37.92 ± 1.36
M2 Y2	3.47 ± 0.09	37.10 ± 1.43
M2 J3	3.46 ± 0.09	37.69 ± 1.39
SM J3	3.44 ± 0.06	38.69 ± 1.33
M1 J1Y1	3.42 ± 0.09	38.26 ± 1.36
M2 J1Y1	3.41 ± 0.07	38.03 ± 1.31
M1 Y2	3.40 ± 0.07	37.14 ± 1.33
M2 SB	3.32 ± 0.07	35.87 ± 1.41
SM Y2	3.27 ± 0.11	38.91 ± 1.34

There are no significant differences in MR on SND and SH of *Prosopis alba* seedlings. However, similar trends to allometric variables (SND and SH) are verified. Thus, the diameter MR shows positive trends with respect to M1 and J3. While the MR of the height of the plantin shows positive trend with respect to M1 and the combination of biostimulant J1 Y1.

4. Conclusions

Our results did not show a clear synergism between the different forms of bioinsumes evaluated considering biochemical and morphological characteristics or by evaluating the mycorrhizal response. Regarding the biostimulants use (regardless of AMF inoculation), foliar application of *L. divaricate* 3% w/v exhibited the lowest values of MDA and total contents of carotenoids. These results confirmed our previous study [11] related to the potential of this biostimulant in enhancing the abiotic stress tolerance of *P. alba*. There were no significant differences, trends indicate positive relationships between neck diameter and mycorrhizal response in *P. alba* seedling treated with M1 with foliar applications of *L. divaricate* at 3% w/v. However, height does not consistently respond to mycorrhizae interacting with biostimulants. Consequently, deeper analysis is needed to understand the effect of the interaction between AMF with biostimulants for improving the plant's physiological status.

Author Contributions: A.C.S.-G., A.T.G., M.G.S., A.d.V.G. and M.B.S. were responsible for conceptualization, methodology, sample collection, analyses, and writing. M.E. assisted with conceptualization, methodology, and resources. C.C., J.G. and R.S.F. participated in sample dating. M.A.N. contributed to the conceptualization, methodology, and writing—review and editing. All authors have read and agreed to the published version of the manuscript.

Funding: The authors are grateful to the projects “Development of a sustainable forestry of implanted forests of high productivity (2019-PE-E1-I016-001)” and “Development and application of technologies for the control of erosion and soil degradation (2019-PE-E2-I052-001) of the Instituto Nacional de Tecnología Agropecuaria (INTA)”. And to the projects “Initial forestry of native species in implanted forests in irrigation zones of Santiago del Estero” and “Biostimulants evaluation to improve the productivity of plant species of regional interest under stress conditions” of the CICYT-UNSE.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data are not publicly available because the data will be used in the future studies.

Acknowledgments: To the Instituto Nacional de Tecnología Agropecuaria (INTA)”, Universidad Nacional de Santiago del Estero (UNSE); Instituto de Fisiología y Recursos Genéticos Vegetales (IFRGV), Centro de Investigaciones Agropecuarias (CIAP), Unidad de Estudios Agropecuarios (UDEA) INTA-CONICET. Estación Experimental Fernández (Universidad Católica de Santiago del Estero-UCSE y Gobierno de la Pcia de Santiago del Estero), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) and the Ministry of Science, Technology and Productive Innovation. Besides, A.C.S.-G. Acknowledges for her fellowship granted by CONICET.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. du Jardin, P. Plant Biostimulants: Definition, Concept, Main Categories and Regulation. *Sci. Hort.* **2015**, *196*, 3–14. [[CrossRef](#)]
2. Ricci, M.; Tilbury, L.; Daridon, B.; Sukalac, K. General Principles to Justify Plant Biostimulant Claims. *Front. Plant Sci.* **2019**, *10*, 494. [[CrossRef](#)] [[PubMed](#)]
3. Carletti, P.; García, A.C.; Silva, C.A.; Merchant, A. Editorial: Towards a Functional Characterization of Plant Biostimulants. *Front. Plant Sci.* **2021**, *12*, 677772. [[CrossRef](#)] [[PubMed](#)]
4. Raymaekers, K.; Ponet, L.; Holtappels, D.; Berckmans, B.; Cammue, B.P.A. Screening for Novel Biocontrol Agents Applicable in Plant Disease Management—A Review. *Biol. Control* **2020**, *144*, 104240. [[CrossRef](#)]
5. Sáenz-Aponte, A.; Correa-Cuadros, J.P.; Rodríguez-Bocanegra, M.X. Foliar Application of Entomopathogenic Nematodes and Fungi for the Management of the Diamond Back Moth in Greenhouse and Field. *Biol. Control* **2020**, *142*, 104163. [[CrossRef](#)]
6. Nguyen, M.L.; Spaepen, S.; du Jardin, P.; Delaplace, P. Biostimulant Effects of Rhizobacteria on Wheat Growth and Nutrient Uptake Depend on Nitrogen Application and Plant Development. *Arch. Agron. Soil Sci.* **2019**, *65*, 58–73. [[CrossRef](#)]
7. Tariq, A.; Pan, K.; Olatunji, O.A.; Graciano, C.; Li, N.; Li, Z.; Song, D.; Sun, F.; Justine, M.F.; Huang, D.; et al. Role of Nitrogen Supplementation in Alleviating Drought-Associated Growth and Metabolic Impairments in Phoebe Zhennan Seedlings. *J. Plant Nutr. Soil Sci.* **2019**, *182*, 586–596. [[CrossRef](#)]

8. Shah, A.; Smith, D.L. Flavonoids in Agriculture: Chemistry and Roles in, Biotic and Abiotic Stress Responses, and Microbial Associations. *Agronomy* **2020**, *10*, 1209. [[CrossRef](#)]
9. Sagadin, M.B.; Monteoliva, M.I.; Luna, C.M.; Cabello, M.N. Diversidad e Infektividad de Hongos Micorrícicos Arbusculares Nativos Provenientes de Algarrobales Del Parque Chaqueño Argentino Con Características Edafoclimáticas Contrastantes. *AgriScientia* **2018**, *35*, 19. [[CrossRef](#)]
10. Salto, C.S.; Sagadin, M.B.; Luna, C.M.; Oberschelp, G.P.J.; Harrant, L.; Cabello, M.N. Interactions between Mineral Fertilization and Arbuscular Mycorrhizal Fungi Improve Nursery Growth and Drought Tolerance of *Prosopis Alba* Seedlings. *Agrofor. Syst.* **2020**, *94*, 103–111. [[CrossRef](#)]
11. Santacruz-García, A.C.; Senilliani, M.G.; Gómez, A.T.; Ewens, M.; Yonny, M.E.; Villalba, G.F.; Nazareno, M.A. Biostimulants as Forest Protection Agents: Do These Products Have an Effect against Abiotic Stress on a Forest Native Species? Aspects to Elucidate Their Action Mechanisms. *For. Ecol. Manag.* **2022**, 522. [[CrossRef](#)]
12. Cavagnaro, T.R.; Smith, F.A.; Ayling, S.M.; Smith, S. Growth and Phosphorus Nutrition of a Paris-type Arbuscular Mycorrhizal Symbiosis. *New Phytol.* **2003**, *157*, 127–134. [[CrossRef](#)] [[PubMed](#)]
13. Giovannini, L.; Palla, M.; Agnolucci, M.; Avio, L.; Sbrana, C.; Turrini, A.; Giovannetti, M. Arbuscular Mycorrhizal Fungi and Associated Microbiota as Plant Biostimulants: Research Strategies for the Selection of the Best Performing Inocula. *Agronomy* **2020**, *10*, 108. [[CrossRef](#)]
14. De Pascale, S.; Roupheal, Y.; Colla, G. Plant Biostimulants: Innovative Tool for Enhancing Plant Nutrition in Organic Farming. *Eur. J. Hortic. Sci.* **2017**, *82*, 277–285. [[CrossRef](#)]
15. Sagadín, M.B. Identificación y Caracterización de los Hongos Micorrícicos Arbusculares Autóctonos en Simbiosis con *Prosopis alba* y los Mecanismos Fisiológicos/Bioquímicos Relacionados con la Tolerancia a Sequía. Ph.D. Thesis, Universidad Nacional de Córdoba, Córdoba, Argentina, 2019.