

Review

The Use of Temperate Tannin Containing Forage Legumes to Improve Sustainability in Forage–Livestock Production

Sebastian P. Lagrange ^{1,*} , Jennifer W. MacAdam ²  and Juan J. Villalba ³

¹ Estación Experimental Agropecuaria Bordenave, Instituto Nacional de Tecnología Agropecuaria, Bordenave, Buenos Aires 8187, Argentina

² Department of Plants, Soils & Climate, College of Agriculture and Applied Sciences, Utah State University, Logan, UT 84322, USA; jennifer.macadam@usu.edu

³ Department of Wildland Resources, Quinney College of Natural Resources, Utah State University, Logan, UT 84322, USA; juan.villalba@usu.edu

* Correspondence: lagrange.sebastian@inta.gob.ar; Tel.: +54-2923-659717

Abstract: Greenhouse gas emissions from ruminant livestock production systems contribute significantly to the environmental footprint of agriculture. Emissions are lower for feedlot systems than for grass-based systems primarily because of the extra time required for grass-finished cattle to reach slaughter weight. In contrast, legume forages are of greater quality than grasses, which enhances intake and food conversion efficiencies, leading to improvements in production and reductions in environmental impacts compared with forage grasses. In addition, the presence of certain bioactives in legumes such as condensed tannins (CT) enhance the efficiency of energy and protein use in ruminants relative to grasses and other feeds and forages. Grazing tannin-containing legumes also reduce the incidence of bloat and improve meat quality. Synergies among nutrients and bioactives when animals graze diverse legume pastures have the potential to enhance these benefits. Thus, a diversity of legumes in feeding systems may lead to more economically, environmentally, and socially sustainable beef production than grass monocultures or feedlot rations.

Keywords: grass-fed beef; sustainable agriculture; forage diversity; tannin-containing legumes; alfalfa; sainfoin; birdsfoot trefoil; condensed tannins; nitrogen excretion; methane emissions



Citation: Lagrange, S.P.; MacAdam, J.W.; Villalba, J.J. The Use of Temperate Tannin Containing Forage Legumes to Improve Sustainability in Forage–Livestock Production. *Agronomy* **2021**, *11*, 2264. <https://doi.org/10.3390/agronomy11112264>

Academic Editor: Agnes van den Pol-van Dasselaar

Received: 12 October 2021
Accepted: 5 November 2021
Published: 9 November 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Emissions of greenhouse gases (GHG) from ruminants include methane (CH₄), nitrous oxide (N₂O), and carbon dioxide (CO₂) [1]. In a recent life cycle assessment of the beef cattle industry in the United States, Rotz et al. [1] estimated that the GHG emissions, considering animal outputs and direct emissions from soil (cultivated pastures, range, and cropland) and the manufacturing of the operation's inputs (fertilizers, pesticides, electricity), were equivalent to 242.6 Tg CO₂eq, which represent 3.8% of the 6457 Tg CO₂eq of total anthropogenic GHG emissions for the US in recent years [2]. Approximately, 142 Tg CO₂eq are directly emitted from cattle systems (CH₄ and N₂O from enteric fermentation and manure management), which is nearly 60% of the total GHG emitted for beef cattle production [1], or 2.1% of the total US anthropogenic GHG emissions [2]. When GHG emissions are expressed per unit of product (GHG intensity), the US average for 2019 was approximately 21 kg CO₂eq/kg carcass weight. In beef production, the cow–calf phase is the biggest contributor, contributing 70% of total GHG emissions [1]. These GHG intensity values are in line with previous values reported by Beauchemin et al. [3] for Canadian beef cattle systems of 22 kg CO₂eq/kg BW, with the cow–calf system contributing approximately 80% of total GHG emissions.

The largest contributing source of GHG emissions from beef cattle production is enteric CH₄, accounting for 56% [1] to 63% [3] of all GHG from beef industry and 39% of all GHG emissions from the livestock sector [4]; thus, reducing emissions from this

source would have the most impact. Methane is a byproduct of the microbial fermentation of feeds in the rumen and may also represent an energy loss to the animal that ranges between 2 to 12% of the gross energy consumed with the diet [5]. The reduction of CO₂ to CH₄ by methanogenic archaea acts as a hydrogen (H₂) sink, removing H₂ from the rumen and avoiding the negative effects of H₂ accumulation on microbial enzymatic activity and degradation of plant material [6]. Methanogens use H₂ as their main energy source, producing CH₄ in the process. Methane is accumulated in the rumen and eructated by the ruminant to the atmosphere [7], resulting in negative implications for the environmental sustainability of ruminant production systems.

Several comprehensive reviews have described different strategies proposed by the scientific community to reduce enteric methane production and mitigate methane emissions [8–12], but in order to be adopted by beef cattle producers, they should be cost-effective and socially acceptable. Rumen defaunation, for instance, has been shown to reduce CH₄ emissions from ruminants by 50%, due to the fact that protozoans are large producers of H₂ with many methanogens among these microorganisms [13]; however, the lack of persistent response due to rapid adaptation and recovery of protozoal numbers along with impractical defaunation methods has limited its use [14]. On the other hand, antimethanogen vaccines have reduced CH₄ emissions up to 8% in sheep [15], but changes in methanogen populations do not always lead to CH₄ reductions [16]. In addition, the development of a successful wide-spectrum immunization is still on the far horizon for CH₄ abatement programs, limiting the application of such strategies as alternatives to reduce CH₄ emissions. Selection of “low-CH₄”-producing animals might also represent a promising strategy for CH₄ mitigation options [17], but the approach is still in an early stage of development. The use of ionophores that inhibit protozoan growth [18], halogenated methane analogues that inhibit growth and enzymatic activity of archaea in the rumen [19], or nitrate salts that have a greater affinity for H₂ than CO₂ [20] have been discouraged due to consumer perception and potential negative effects on animal health, as well as on the environment [21].

Finally, dietary manipulations such as feeding highly digestible feed components such as grains [22] or feeding organic acids such as fumarate [23] or malate [24], which promote propionate production in the rumen and redirect H₂ to other reductive bacteria, may reduce CH₄ emissions from ruminants. The addition of lipids [25], condensed tannin (CT) extracts [26,27], essential oils [28], exogenous enzymes and yeasts [29], among others—which can be supplied through total mixed rations for confined livestock—are still the most promising CH₄ mitigation options in terms of practical application and acceptance by farmers and consumers. Nevertheless, many ruminants consume forages as their sole diet in pasture-based livestock systems, and the need to supply feed additives in meals might constrain their practical implementation [21]. In this case, CH₄ emissions may be reduced by using highly digestible forage species with a low concentration of fiber [30,31], because forages such as grasses with high fiber concentration reduce passage rate and increase ruminal retention time [32,33], which enhances CH₄ production per unit of forage intake (CH₄ yield). In this situation, the extent of rumen fermentation increases and there is more H₂ to be used as a substrate for methanogenic archaea [34]. In addition, a more fibrous diet usually increases the proportion of acetate to propionate in the rumen, which increases the production and release of CH₄ [5,35].

2. The Use of Forage Legumes for Enteric Methane Abatement in Forage-Based Beef Production Systems

Forage legumes in beef feeding systems can offer economic and environmental advantages relative to grass-fed systems. In contrast to grasses, forage legumes are lower in neutral detergent fiber (NDF), higher in N concentrations [36,37], higher in nonstructural carbohydrates [38,39], and are digested more rapidly by ruminants at similar stages of maturity [37]. These characteristics lead to lower retention times in the rumen; thus, intake and production are greater than in grass-fed systems [40]. This faster rate of digestion of forage legumes is primarily attributed to the faster rates of particle breakdown and

quicker fermentation rates in the rumen [41]. An increased passage rate of forage legumes may also favor propionate production, which is considered a competitive pathway for H₂ use in the rumen [34], contributing to reduced CH₄ yield relative to other forages such as grasses [42]. In support of this, Archimède et al. [43], in a meta-analysis of ruminants fed C₃ or C₄ grasses and legumes, identified fiber structure and ruminal retention time as the main factors influencing CH₄ production, with 20% lower CH₄ yields in animals fed warm-season legumes than in those fed C₄ grasses. Similarly, the enteric CH₄ emissions of beef cows grazing the forage legume birdsfoot trefoil (*c*) and cicer milkvetch (*Astragalus cicer*) were approximately half of the emissions reported for the grass meadow brome (*Bromus biebersteinii*) [44].

Alternatively, forages with high concentrations of nonfiber carbohydrates (NFC; soluble carbohydrates plus pectin) that are rapidly fermented in the rumen, and with a low proportion of structural carbohydrates (cellulose and hemicellulose), may yield levels of microbial mass similar to those observed in grain-fed animals, increasing proportions of potentially propionate-forming bacteria and reducing H₂ production and CH₄ emissions, as was observed by Sun et al. [42] feeding forage rape (*Brassica napus* L.) to lambs. Nonfiber carbohydrates represent a readily fermentable source of energy for microorganisms in the rumen, providing energy in synchrony with the high concentrations of protein availability typically observed in forage legumes that contribute to the synthesis of microbial protein [45].

The high nutritional composition of legumes usually leads to greater DM intakes than in animals offered grasses [37], resulting in greater liveweight gains (0.8 to 1.6 kg/d for beef steers) [44,46–48]. For finishing animals, this benefit substantially decreases the number of days to slaughter and the amount of CH₄ emitted over the animal's lifetime relative to grass-fed systems [37]. It has been estimated that the number of cattle required to produce 1 billion kg of beef when finished on pure birdsfoot trefoil pastures is approximately 15% lower than when finished on grass (2.9 vs. 3.4 million animals, respectively; [48]), suggesting that legume-finishing systems represent a realistic strategy to reduce enteric CH₄ emissions.

3. Constraints to the Use of Forage Legumes in Beef Cattle Grazing Systems

Alfalfa (*Medicago sativa* L.) has been one of the most important crops grown in the western US, being the most high-yielding and nutritious forage available for feeding high-producing ruminants [49]. Similarly, white clover (*Trifolium repens*) and red clover (*T. pratense*) have been extensively used for grazing in Australia, New Zealand, and the United Kingdom. However, the direct use of these legumes as grazing forage has been limited due to the high risk of livestock losses caused by pasture bloat [50]. Pasture bloat occurs when ruminants graze fresh, high-protein forages with a high rate of particle breakdown that results in rapid release of plant-soluble proteins and disruption of chloroplasts, providing large quantities of gas and bacterial slime, which create a stable foam that prevents eructation of fermentation gases (CO₂ and CH₄) [51]. Ultimately, the rumen becomes distended, resulting in death from suffocation or cardiac arrest. Subclinical bloat is another significant but often unnoticed cause of reductions in productivity, mostly explained through reductions of intake [52]. Management techniques such as grazing mature bloat-causing legumes might reduce the risk at the expense of reducing the overall nutritive value of legume forages [53]. Grazing grass–legume mixtures still may impose a risk of bloat if animals are able to select and ingest the preferred legume species in high proportions.

A further issue with alfalfa and *Trifolium* spp. is that the high concentration of rumen-degradable protein in forage legumes usually exceeds the capacity of microorganisms for uptake of NH₃ and synthesis of microbial protein due to a deficient energy supply for N capture [54]. The excess of ruminal NH₃ is absorbed across the rumen wall [55], transformed to urea in the liver, and excreted in the urine with an energy cost for the animal [56]. Consequently, only 10 to 40% of ingested N is retained as animal product

(meat or milk) by ruminants [57], and in some cases, when NH_3 detoxification capacity of the liver is surpassed, NH_3 accumulation in blood can be toxic for the ruminant and induce negative internal states that constrain DM intake [58]. In addition, high blood urea levels lead to high urinary N excretions [59] that increase the proportion of N excreted as a highly labile form in the urine, contributing to pollution from agricultural sources that is a major environmental concern [60].

Once urine is excreted and deposited on the soil surface, urea is rapidly hydrolyzed by microbial urease to NH_4^+ , which may be nitrified later to nitrite (NO_2^-) and nitrate (NO_3^-) [61]. Greater levels of urinary N excretions are associated with a greater and more rapid NH_3 volatilization and N losses as NO_3^- that may be leached into groundwater or in runoff to waterways [60], contributing to eutrophication [62,63] and the pollution of drinking water. In addition, nitrous oxide (N_2O) is produced as an obligate intermediary during microbial nitrification and denitrification processes [64,65], being one of the most important GHGs with a warming potential 265 times greater than CO_2 in a 100-year time horizon [66]. According to Bao et al. [67], an increment in urinary N excretion of growing beef cattle from 29 to 50 g/d increases the estimated emission of N_2O by 37% from 413 to 565 mg/d. Regardless of these conditions, reductions in the proportion of N partitioned to urine in ruminants will be beneficial for the environment, since urinary N is much more susceptible to gaseous losses than fecal N, which is in the form of covalently bound N and needs a longer time to be mineralized to NH_4^+ before being susceptible to volatilization or available for nitrification [68].

4. Tannin-Containing Legumes in Forage-Based Livestock Systems

To counteract the high urinary N excretion that can result from the grazing of some temperate legumes, legume species that contain moderate concentrations (i.e., 30–60 g/kg DM basis) of the bioactive secondary compounds CT, including sainfoin (*Onobrychis viciifolia*) and birdsfoot trefoil (*Lotus corniculatus*), are used. Sainfoin is a legume species that naturally contains significant concentrations of CT (30 to 80 g CT/kg DM; [69]) distributed throughout the aerial parts of the plant and restricted to the cell's vacuoles [70]. Sainfoin can serve either as an alternative or associate forage crop to alfalfa pastures in climate-adapted environments. The yield and nutritive value of sainfoin are comparable to alfalfa [71], leading to similar performance of sheep and cattle [72–74]. In fact, heifers grazing a 3-way choice among sainfoin, birdsfoot trefoil, and alfalfa selected a varied diet, preferring sainfoin over birdsfoot trefoil or alfalfa in a 46:27:27 ratio, and in a 70:30 ratio when cattle could choose between sainfoin–birdsfoot trefoil or sainfoin–alfalfa, respectively [75].

Birdsfoot trefoil, on the other hand, is a legume species that presents a more prostrate growth habit relative to alfalfa or sainfoin [76], with greater biomass per unit of area and higher bulk density (i.e., herbage weight per unit of canopy volume), which is correlated with a greater leaf area index [77]. It contains 10 to 40 g CT/kg DM [78] and yields approximately two-thirds as much as alfalfa in pure stands in the northern Mountain West [79], with a nutritional value similar to alfalfa [76]. Thus, the use of tanniferous legumes in monocultures or associated with other nontanniferous legumes may reduce ruminal protein degradability and alleviate malaise by inhibiting NH_3 production in the rumen. This strategy will increase the pool of high-quality protein that reaches the small intestine [80], shifting N excretion from urine to feces while improving N utilization [81].

4.1. Condensed Tannin Structure

Condensed tannins are plant secondary compounds (PSCs) also known as proanthocyanidins, consisting of oligomers or polymers of flavan-3-ol monomers that differ due to the hydroxyl groups and the stereochemistry (spatial orientation) of C-2 and C-3 in the C-ring [82]. Most of the CT occurring in forage species are procyanidin (PC) (e.g., catechin and epicatechin) and prodelphinidin (PD) subunits (e.g., galocatechin and epigallocatechin), which possess an additional hydroxyl group at C-5 of the B-ring [83]. Epicatechin

and epigallocatechin have a cisorientation of the C-2 and C-3 in the C-ring, while catechin and gallicocatechin possess a transorientation (see Zeller, [83]).

Monomers grow into oligomers and polymers through covalent linkages of the C-4 in the C-ring of a flavan-3-ol to the C-8 or C-6 positions in the C-ring of another monomer [84] (Figure 1). These oligomers and polymers in common forage plants are typically present as mixtures of PC and PD subunits, which are distributed throughout the CT molecule, linked at different positions, leading to many different chemical structures within CT [83]. Molecules of CT also differ in the number of flavan-3-ol subunits they contain (degree of polymerization), resulting in structures that can vary in MW between 1900 and 28,000 Da [82]. Thus, plants' CT vary in degrees of polymerization and the composition of their subunits, and they can differ among plant species, cultivars within the same species, and even organs (leaves, stems, roots) within the same plant [85]. In addition, the concentration of CT varies with phenological stage, declining in concentration as maturity progresses [86]. For instance, leaves of sainfoin have higher CT concentrations and a greater biological activity and PD proportion than stems [87]; therefore, vegetative stages contain higher concentration of CT than mature plants [88], and thus, a greater CT-protein complexation potential [89].




Figure 1. Condensed tannin molecule consisting of four flavan-3-ol monomers. Reproduced with permission from Mueller-Harvey et al. [84], Crop Science; published by John Wiley & Sons, 2019.

4.2. Condensed Tannin-Protein Complexes and Reductions in Urinary N Excretions

Once plant tissues are chewed or degraded during microbial digestion, CT are released from vacuoles and bind to plant salivary and microbial proteins forming insoluble complexes in the rumen [90]. These complexes reduce protein solubilization and protect dietary proteins from microbial hydrolysis and deamination in the rumen, reducing the susceptibility of forage protein to microbial degradation [91]. In addition, CT can form complexes with extracellular and cell coat enzymes of proteolytic bacteria, inhibiting their activity and reducing protein degradation [92]. As a result, there is an increased outflow of undegraded plant protein to the intestines, and reductions in ruminal NH_3 concentrations [93–95]. The CT-protein complexes are stable over the pH range from 3.5 to 7.0 but can dissociate in the abomasum and anterior duodenum at a lower pH [96], releasing proteins for gastric and peptic digestion and increasing the proportion of plant amino acids available for postruminal absorption [97], increasing the efficiency of N utilization by the ruminant.

The formation of the CT-protein complex is due to hydrogen bonding interactions between the hydroxyl groups ($-\text{OH}$) of the CT molecule and the amino group ($-\text{NH}_2$) of peptides (Figure 2), or by hydrophobic interactions between the phenol ring and the carboxyl group ($-\text{COOH}$) of proteins [90]. These are weak associations involving noncovalent CT-macromolecule interactions. The formation of such complexes depends on the structure

of both the protein and the specific CT in the plant or plant part, the isoelectric point of the protein, the pH in the gastrointestinal tract, and the tannin–protein molar ratios [85].




Figure 2. Hydrogen bonding involved in condensed tannin–protein complexation. Reproduced with permission from Zeller [83], Crop Science; published by John Wiley & Sons, 2019.

Different studies have determined that as CT concentration or MW and mean degree of polymerization increase, the protein precipitation capacity of CT also increases [98,99]. Aufrère et al. [100] found a negative correlation between N solubility and CT concentration, PD/PC ratio, mean degree of polymerization, and cis/trans ratio for three sainfoin varieties at several harvests.

Biochemical mechanisms of bonding between polyphenols and macromolecules also involve irreversible covalent interactions mediated by oxidation of phenolic compounds with the formation of o-quinones or o-semi-quinones, or through the cleavage of proanthocyanidin bonds with the formation of carbocations [101]. Covalent interactions have received less attention than noncovalent interactions, although they have been demonstrated between polyphenols and individual amino-acids [101].

Condensed tannins in birdsfoot trefoil have average molecular weights of 4400 Da [102], with a degree of polymerization in the range of 6 to 14 of predominantly PC type subunits [90], while sainfoin's CTs are predominantly constituted of PD monomers with a mean MW of 5100 Da [102], with polymer sizes that vary between 4–12 subunits [90]. Thus, differences between the molecular structure of CT between birdsfoot trefoil and sainfoin may result in different effects on protein degradability because they differ in binding capacities and affinities for plant, microbial, and mammalian proteins during herbivory. This may explain the higher protein precipitation capacity reported for sainfoin's CT relative to CT from birdsfoot trefoil [102].

Sainfoin has been found to decrease urinary N losses by ruminants [87,103]. Several *in vitro* [104] and *in vivo* studies [87,105,106] have reported reductions in ruminal protein degradation, ruminal NH₃ concentrations, and urinary N excretion of substrates incubated with sainfoin or of sheep fed sainfoin relative to animals receiving polyethylene glycol (PEG), a polymer that binds to CT more readily than protein [107]. Condensed tannins in sainfoin may also enhance ruminant nutrition relative to other perennial legumes such as alfalfa [69]. In an *in vitro* study, Williams et al. [108], found that NH₃ concentrations were lower when sainfoin was incubated in continuous cultures than when alfalfa (a nontanniferous legume) was used as the substrate. However, NH₃ was not different between birdsfoot trefoil and alfalfa in this study. Similar results were obtained later by

Grosse Brinkhaus et al. [109], who observed a 21% reduction in blood urea N and a 38% lower urinary N when dairy cows were fed sainfoin than when they were fed alfalfa pellets; and Lagrange et al. [74] reported that yearlings heifers consuming sainfoin or birdsfoot trefoil showed a 40% reduction in urinary N concentration relative to those grazing alfalfa, diverting more of the N to feces, thereby reducing the loss of N as ammonia into the atmosphere. This study also demonstrated that the partial replacement of alfalfa by sainfoin and birdsfoot trefoil in 2-way or 3-way choices was also effective in reducing the urinary N concentrations of beef heifers. Similarly, Aufrère et al. [110] showed in vitro that mixing sainfoin with alfalfa could be an efficient way to reduce the N solubility of pure alfalfa. Finally, tannin-containing hays have also shown potential to reduce urinary urea N excretion, increase N retention, and reduce enteric CH₄ emissions from beef cattle, suggesting that CTs remain active during the process of forage conservation [111]. Thus, ecoregions around the world where legume supply is limited during certain times of the year, such as spring or winter, could benefit from the provision of CTs through the use of preserved forages.

However, when sainfoin is fed to ruminants, CT-protein complexes may not be completely dissociated in the abomasum and continue intact through the small intestine, preventing amino acid digestion and absorption [97,112]. The potential of these complexes for being reversible is dependent on the type of bonding (noncovalent or covalent) between CT and proteins [101]. Alternatively, CT may still be active under the pH level (5.0) of the proximal small intestine and interfere with endogenous and microbial proteolytic enzymes, increasing the proportion of protein in the feces [94]. In support of this, Lagrange et al. [74] observed that beef cattle grazing sainfoin partitioned more N to feces (30.1% vs. 22.7%, respectively) than animals grazing birdsfoot trefoil, and sheep fed fresh sainfoin showed greater fecal N than sheep fed pure birdsfoot trefoil or alfalfa (31.5% vs. 26.6%, respectively; [113]). This may reduce N retention, as observed for sainfoin diets [114].

The prevalence of PC subunits in birdsfoot trefoil tannin may be associated with a greater protein digestion in the abomasum and small intestine and improved amino acid absorption [81,90]. A greater amino acid absorption has been linked to overall improvements in animal performance, including body weight gain, wool and milk production, reproductive performance, and the ability to cope with gastrointestinal nematode burdens [115]. For instance, Min et al. [116] reported increments of reproduction efficiency and wool production in sheep fed birdsfoot trefoil relative to animals receiving PEG, a polymer that binds and inactivates tannins. This response was produced without increments in voluntary intake, but authors reported a greater concentration of plasma-essential amino acids, suggesting a higher intestinal absorption. The unique CT produced by birdsfoot trefoil, as well as its high fiber digestibility [117–119], also enhance the efficiency of energy and protein use in ruminants relative to other nontanniferous legumes. Sheep grazing birdsfoot trefoil had significantly improved performance compared with sheep grazing alfalfa pastures, resulting in greater ewe and lamb weight gains, carcass dressing-out percentage, and wool growth [120]. Harris et al. [121] found that dairy cows grazing birdsfoot trefoil improved the efficiency of feed utilization and increased milk yield by 10%, with increments in milk protein concentration relative to white clover (another nontanniferous legume), and Lagrange et al. [74] reported 40% greater average daily gains (ADG) in beef heifers grazing birdsfoot trefoil relative to animals grazing alfalfa.

4.3. Effect of Condensed Tannins on Enteric Methane Emissions

Condensed tannins may inhibit CH₄ production in the rumen, which is beneficial for improving nutrient utilization and reducing dietary energy loss and GHG emissions for eco-friendly animal production. Several studies have reported reductions (13–16%) either in the gross emission of CH₄ (g/d) or in CH₄ yield (g/kg dry matter intake), using forages with moderate concentrations of CT (20–50 g/kg DM) [122–124], or CT-containing plant extracts supplied with the feed [125] or drenched directly to the animals [27]. A meta-analysis from

15 *in vivo* experiments showed that increasing tannin concentration in the diet decreased CH₄ production linearly when expressed relative to dry matter intake (DMI) or digestible OM intake [126]. Thus, low concentrations of CT (<20 g/kg DM) may not affect CH₄ production in ruminants relative to control diets [82].

Chemical structure of CT may also be an important factor affecting enteric CH₄ production, as was demonstrated *in vitro* by Hatew et al. [127], who found differences in CH₄ emissions among CT extracts from four different sainfoin accessions. Sainfoin is a legume species that has been shown to reduce CH₄ production in *in vitro* studies [128–130]. As mentioned previously for the protein precipitation capacity of CT, as the degree of polymerization of CT increases, greater reductions in CH₄ production have been reported for *in vitro* studies [131]. Likewise, higher molecular weight fractions of CT significantly decreased total methanogen numbers *in vitro* compared with lower molecular weight CT fractions [132].

The effect of CT on enteric CH₄ emissions has been attributed to a direct effect on methanogenic archaea and/or their enzymatic activity [131–133], or more likely, to an indirect effect on fiber digestion, adversely affecting cellulolytic bacteria and consequently reducing the amount of forage substrate fermented in the rumen (reduced digestion) [126,134,135]—a process that may be subsequently compensated in the lower digestive tract by colonic fermentation [40]. The bacteria that digest cellulose produce both acetate and H₂. However, accumulation of H₂ inhibits fermentation, so Archaea dispose of H₂ by using it to reduce CO₂ to CH₄ [34]. Condensed tannins likely inactivate extracellular microbial enzymes through the formation of CT–enzyme complexes, subsequently reduce their digestive activity [136], and/or directly inhibit cellulolytic bacteria [137]. In addition, formation of cell-associated protein–tannin complexes on the cell surface may interfere with microbial attachment to fiber and prevent microbial digestion [138]. In support of this, Wang et al. [69] and Barry and McNabb suggested that concentrations of CT in forages greater than 50 g/kg might decrease DM digestibility in ruminants, and Chung et al. [139] observed a lower NDF digestibility in sainfoin than in alfalfa (45.3 vs. 55.3%), even with CT concentration in sainfoin as low as 2.45%. Reduced fiber digestion due to an increased CT ingestion may also slow clearance of forage residues from the rumen, reducing voluntary DMI [81]; thus, reductions in enteric CH₄ emissions due to a decreased fiber digestibility would not be a viable strategy.

Ciliate protozoa also produce H₂; reducing their numbers with rumen defaunation by supplying CT with the ingestion of tropical legumes [140] could indirectly affect CH₄ emissions, as mentioned previously, either by reducing methanogens symbiotically associated with protozoal populations or by reducing fiber digestion and H₂ supply to methanogenic archaea [141].

Rumen microbiome adaptation to plant secondary compounds is possible, which could influence long-term bioactivity and, thus, enteric CH₄ production, although information on this topic is still limited [142,143]. Such adaptation may be influenced by the specific chemical structure of the phenolic compound in question. For instance, rumen microbes have been reported to adapt to chemicals such as carvacrol and thymol to a greater extent than to phenolics in garlic oil [144]. Further research is needed to determine if the duration of feeding tannin-containing legumes influences the rumen microbiome and if methanogenic adaptation occurs.

5. Other Beneficial Effects of Tanniferous Legumes in Grazing Beef Production Systems

Another advantage of grazing tanniferous legumes is a reduction of the risk of bloat [50], which allows cattle to graze forage legumes at the greatest nutritional value. Tanniferous legumes such as birdsfoot trefoil and sainfoin are nonbloating and can therefore be grazed in pure stands. Complexes between CT and proteins prevent the plant protein from being solubilized into ruminal fluid, inhibiting the formation of proteinaceous, gas-trapping foam [145]. It has been calculated that CT concentrations as little as 1 to 5 g/kg DM should prevent bloat [146]. Adding a source of CT to highly digestible alfalfa could

reduce the availability of soluble protein and the rate of gas production and proliferation of ruminal microbial populations, preventing the formation of persistent foam [50]. In support of this, the inclusion of sainfoin into alfalfa pastures has reduced the incidence of bloat [147] and may therefore be a practical and effective means of controlling this disorder. McMahon et al. [128] reported a marked reduction in pasture bloat when as little as 10% sainfoin was included in fresh alfalfa diets.

In addition to their positive effects at attenuating bloat and reducing environmental impacts, the use of forage legumes for finishing beef cattle also results in greater carcass weight, dressing percentage, backfat thickness, and intramuscular fat percentage in the longissimus muscle compared with grass-finished beef (4.4% vs. 2.9%, respectively), approaching values observed for grain-based finishing systems (5.8%; [148]). This outcome might be related to the high concentration of NFC present in forage legumes. Likewise, tenderness, fattiness, juiciness, and overall liking of legume-finished beef did not differ from grain-finished beef, and both types of beef presented greater scores for these characteristics than grass-fed beef [148]. In addition to these results, the omega-6 to omega-3 fatty acid ratio observed in legume-finished beef was much lower than that observed with concentrate diets and similar to grass-fed diets (2.41, 5.74, and 3.44, respectively), with greater omega-3 as well as reduced omega-6 in legume-finished beef [148], maintaining the benefits for human health of the consumption of healthy fatty acids [149–151]. Furthermore, as CTs reduce the activity of specific rumen bacteria responsible for biohydrogenation of dietary fatty acids [152], tanniferous legumes such as sainfoin may promote increments in conjugated linoleic acid and polyunsaturated fatty acids and reductions in saturated fatty acids in meat relative to animals consuming diets without CT [153]. In support of this, beef carcasses from cattle that were fed sainfoin had greater marbling scores, quality grades, and backfat thicknesses than alfalfa-fed cattle, and steaks were redder in color than steaks from cattle finished on alfalfa and contained more unsaturated fatty acids [73].

In contrast to both cereal grains and pasture grasses, perennial legumes form symbiotic associations with soil bacteria (*Rhizobia* spp.) and fix their own N, being productive for multiple years without the need of N fertilization [154,155]. Finishing cattle on N-fixing forages reduces costs and enhances ranch profitability while reducing environmental impact associated with the use of N fertilizers, as they increase nitrate production and water eutrophication [61–63]. Additionally, GHG emissions related to the production, transport, and use of N-based fertilizers [37] decrease in forage legume systems. Moreover, direct emissions of nitrous oxide (N₂O) are negligible from biological N fixation [156].

Therefore, legume-fed beef production systems give producers a sustainable alternative forage-based livestock production program while maintaining high animal performance and high-quality beef that is comparable to grain-finishing programs. In addition, the use of tannin-containing legumes either as pure forages or in association with CT-free legumes offer a feasible solution to the problems of low N utilization and high risk of bloat for cattle grazing nontanniferous legume monocultures, increasing the efficiency of N use and improving the health of ruminants, humans, and the environment.

6. Forage Diversity in Beef Cattle Production Systems

A diversity of forages and biochemicals available in pasturelands may enhance the benefits described above because the complementary relationships among multiple food resources in nature improves the fitness of herbivores [157], which in turn, can reduce environmental impacts. Herbivores have evolved grazing in diverse plant communities, consuming arrays of feeds with different chemical and physical characteristics [158]. Diverse diets offer ruminants a variety of nutrients and PSCs, which allow for a more balanced diet with greater medicinal benefits than single forage species in monocultures [159,160]. In addition, complementarities among nutrients and PSCs may lead to more efficient use of feeds, with improvements in animal welfare and productivity [161], and reduced carbon and N emissions to the environment [115,162]. For instance, the consumption of different forage species with contrasting chemical compositions (different concentrations of NFC,

fiber, and proteins) and the presence of CT may lead to associative effects, such as protein degradability lower than the average of the individual forages, as it has been demonstrated in *in vitro* conditions by Niderkorn et al. [163] for a mixture of sainfoin and cocksfoot (*Dactylis glomerata*).

Some bioactive secondary metabolites in forage legumes can cause digestive interactions, so that the rumen fermentation pattern of a mixture of forages can differ from the average values of its components [164], resulting in positive (synergistic) or negative (antagonistic) effects on ruminant nutrition. It may therefore be preferable to use more than one CT source so that individual sources are ingested at a lower dosage to avoid potential antinutritional effects of high concentrations of single CT [164]. As described previously, tannins produced by different forage species, cultivars, plants, plant parts, or during different seasons may have contrasting physical and chemical properties that may impact herbivores in different ways [161]. Thus, mixtures between legumes with different CT chemical structures may produce associative effects that enhance the effect relative to a single CT. This was demonstrated by Lagrange et al. [74] in an *in vivo* study where yearling heifers grazing a choice of tanniferous legumes (birdsfoot trefoil and sainfoin) showed lower levels of urinary N concentration than animals grazing the same legumes in monoculture, suggesting a synergism between different CT sources. In addition, this study showed that heifers grazing the tanniferous legumes (birdsfoot trefoil and sainfoin) in association with a non-tannin-containing legume (alfalfa) reduced urinary N excretion (40.7 vs. 50.6%) and retained more N (36.1 vs. 25.2%) relative to control animals grazing the same species as monocultures on average, respectively [74]. Previously, Aufrère et al. [165] had demonstrated that CT from sainfoin could bind and precipitate protein from alfalfa.

Interactions among CT may also influence the total amount of food a herbivore can ingest [166,167]. It has been observed that the DMI of sheep increases as the number of tanniferous shrubs in the diet increases, relative to single-shrub diets [167]. Food diversity may also provide ruminants a positive stimulus that increases their motivation to eat [168]. A diversity of forages allows animals to incorporate different species into their diets, which may delay the onset of satiety [169]. In contrast, animals constrained to monocultures may reach satiety at lower levels of feed intake due to the nutritional imbalances or excessive orosensory exposure to limited stimuli. In support of this, Lagrange et al. [74] observed a 33% greater dry matter intake (DMI) and 30 to 50% greater ADG in heifers that grazed a 3-way choice among alfalfa, sainfoin, and birdsfoot trefoil relative to the average DMI and ADG from animals grazing the same three species in monocultures, suggesting a synergism among pasture species when these were consumed together. Similarly, sheep that were offered a choice of different legumes and selected 50% alfalfa, 35% sainfoin, and 15% birdsfoot trefoil had 10% greater DMI and DM digestibility than for the average DMI value calculated from the same proportions of these forages when they were fed separately [113]. On the other hand, Wang et al. [147] observed similar feed intakes in beef steers grazing pure alfalfa or mixed alfalfa–sainfoin pastures containing up to 35% sainfoin; Christensen [170], when feeding a mixture of alfalfa–birdsfoot trefoil hays to dairy cows, did not find differences in DMI relative to feeding pure alfalfa.

Animals that are motivated to eat different species (i.e., a choice of legumes) could also incur greater energy expenditures in order to gather different forages and achieve the challenge of building a balanced diet [171]. The spatial aggregation of forage species in contiguous swards as opposed to an intermingled mixture may reduce search time allowing animals being more efficient in diet selection [172]. In a finely intermingled mixed pasture, animals may have a reduced intake rate due to time spent searching for the preferred plant species [173] and reduces daily voluntary intake relative to grazing monocultures. Moreover, some less competitive species such as sainfoin may be outcompeted in a mixture with better adapted species such as alfalfa, or the most preferred herbage species could be overgrazed, leading to resource degradation [174,175]. In contrast, when three different forage legumes were established in side-by-side patches, beef heifers grazing the choice

treatments did not invest extra time in walking, searching, or patch switching activities relative to heifers grazing monocultures [75].

Finally, giving choices to ruminants and allowing them to solve problems of nutrient imbalances or excess exposure to a single PSC may elicit positive emotional states and ultimately improve their welfare relative to animals limited to monocultures [176]. Animals exposed to a diverse array of foods have lower indicators of stress relative to animals ingesting single rations [177] and have the opportunity to learn the postingestive consequences of foods and how to meet their needs through selecting a varied diet [178]. Diversity also allows animals to select a diet that is a function of their specific and dynamic needs. In contrast, rations designed for the “average” individual may not satisfy all animals’ needs given the inherent individual differences that exist among animals [179].

7. Conclusions

Today’s beef producers are challenged by growing consumer demand for environmentally, economically, and socially sustainable food [180,181], and consumer attention to production sources, animal welfare, and human health is on the rise [182,183]. In the US and other beef-producing countries, wetlands and grasslands have been converted to cropland that is used for the production of annual cereal grains, the majority of which are fed to livestock rather than consumed by humans. The ecosystem services of annually cropped farmland are compromised by reduced organic matter, periods of bare soil, and frequent application of inorganic soil nutrients. At the other extreme, grass-finished production systems provide a food source of relatively low quality that reduces productivity, increasing time to slaughter and related environmental impacts. In contrast to grasses, perennial legumes fix their own nitrogen and are digested more rapidly than grasses by ruminants; thus, the intake, production, and efficiency of feed conversion to red meat or milk are higher than for forage grasses, resulting in reduced environmental impacts compared with grass-fed cattle. The unique tannins produced by some legumes such as birdsfoot trefoil and sainfoin, as well as the high fiber digestibility of temperate forage legumes, enhance the efficiency of energy and protein use in ruminants relative to non-tannin-containing legumes such as alfalfa. Synergisms achieved by a diversity of legumes with beneficial PSCs may further enhance the benefits observed for single species, contributing to the development of beef production systems that improve overall sustainability with reduced environmental impacts while satisfying human food needs.

Author Contributions: Conceptualization, S.P.L., J.W.M. and J.J.V.; writing—original draft preparation, S.P.L.; writing—review and editing, J.J.V. and J.W.M. All authors have read and agreed to the published version of the manuscript.

Funding: This effort was supported by the Utah Agricultural Experiment Station (grant number UTA0 1321) and the National Institute of Food and Agriculture (NIFA), USDA (award no. 2016-67019-25086). This paper was published with the approval of the Director, Utah Agricultural Experiment Station, and Utah State University, as journal paper number 9524.

Institutional Review Board Statement: Not Applicable.

Informed Consent Statement: Not Applicable.

Data Availability Statement: Not Applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Rotz, C.A.; Asem-Hiablie, S.; Place, S.; Thoma, G. Environmental Footprints of Beef Cattle Production in the United States. *Agric. Syst.* **2019**, *169*, 1–13. [CrossRef]
2. EPA. Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990–2017. Available online: <https://www.epa.gov/ghgemissions/inventory-us-greenhouse-gas-emissions-and-sinks-1990-2017> (accessed on 24 January 2020).
3. Beauchemin, K.A.; Henry Janzen, H.; Little, S.M.; McAllister, T.A.; McGinn, S.M. Life Cycle Assessment of Greenhouse Gas Emissions from Beef Production in Western Canada: A Case Study. *Agric. Syst.* **2010**, *103*, 371–379. [CrossRef]

4. Gerber, P.J. *Tackling Climate Change through Livestock: A Global Assessment of Emissions and Mitigation Opportunities*; Food and Agriculture Organization of the United Nations, Ed.; Food and Agriculture Organization of the United Nations: Rome, Italy, 2013; ISBN 978-92-5-107920-1.
5. Johnson, K.A.; Johnson, D.E. Methane Emissions from Cattle. *J. Anim. Sci.* **1995**, *73*, 2483–2492. [[CrossRef](#)]
6. McAllister, T.A.; Newbold, C.J. Redirecting Rumen Fermentation to Reduce Methanogenesis. *Aust. J. Exp. Agric.* **2008**, *48*, 7. [[CrossRef](#)]
7. Janssen, P.H. Influence of Hydrogen on Rumen Methane Formation and Fermentation Balances through Microbial Growth Kinetics and Fermentation Thermodynamics. *Anim. Feed Sci. Technol.* **2010**, *160*, 1–22. [[CrossRef](#)]
8. Broucek, J. Options to Methane Production Abatement in Ruminants: A review. *J. Anim. Plant. Sci.* **2018**, *28*, 348–364.
9. Haque, M.N. Dietary Manipulation: A Sustainable Way to Mitigate Methane Emissions from Ruminants. *J. Anim. Sci. Technol.* **2018**, *60*, 15. [[CrossRef](#)]
10. Alemneh, T.; Getabalew, M. Strategies to Reduce Methane Emission in Ruminants. *Int. J. Ecol. Ecosolution* **2019**, *6*, 16–22.
11. Pámanes-Carrasco, G.; Herrera-Torres, E.; Murillo-Ortiz, M.; Reyes-Jáques, D. Climate Change Mitigation in Livestock Production: Nonconventional Feedstuffs and Alternative Additives. In *Livestock Health and Farming*; IntechOpen: London, UK, 2019. [[CrossRef](#)]
12. Islam, M.; Lee, S.-S. Advanced Estimation and Mitigation Strategies: A Cumulative Approach to Enteric Methane Abatement from Ruminants. *J. Anim. Sci. Technol.* **2019**, *61*, 122–137. [[CrossRef](#)]
13. Hegarty, R.S. Reducing Rumen Methane Emissions through Elimination of Rumen Protozoa. *Aust. J. Agric. Res.* **1999**, *50*, 1321. [[CrossRef](#)]
14. Martin, C.; Morgavi, D.P.; Doreau, M. Methane Mitigation in Ruminants: From Microbe to the Farm Scale. *Animal* **2010**, *4*, 351–365. [[CrossRef](#)]
15. Wright, A. Reducing Methane Emissions in Sheep by Immunization against Rumen Methanogens. *Vaccine* **2004**, *22*, 3976–3985. [[CrossRef](#)] [[PubMed](#)]
16. Williams, Y.J.; Popovski, S.; Rea, S.M.; Skillman, L.C.; Toovey, A.F.; Northwood, K.S.; Wright, A.-D.G. A Vaccine against Rumen Methanogens Can Alter the Composition of Archaeal Populations. *Appl. Environ. Microbiol.* **2009**, *75*, 1860–1866. [[CrossRef](#)] [[PubMed](#)]
17. Pickering, N.K.; Oddy, V.H.; Basarab, J.; Cammack, K.; Hayes, B.; Hegarty, R.S.; Lassen, J.; McEwan, J.C.; Miller, S.; Pinares-Patiño, C.S. Animal Board Invited Review: Genetic Possibilities to Reduce Enteric Methane Emissions from Ruminants. *Animal* **2015**, *9*, 1431–1440. [[CrossRef](#)] [[PubMed](#)]
18. Guan, H.; Wittenberg, K.M.; Ominski, K.H.; Krause, D.O. Efficacy of Ionophores in Cattle Diets for Mitigation of Enteric Methane. *J. Anim. Sci.* **2006**, *84*, 1896–1906. [[CrossRef](#)] [[PubMed](#)]
19. Goel, G.; Makkar, H.P.S.; Becker, K. Inhibition of Methanogens by Bromochloromethane: Effects on Microbial Communities and Rumen Fermentation Using Batch and Continuous Fermentations. *Br. J. Nutr.* **2009**, *101*, 1484. [[CrossRef](#)] [[PubMed](#)]
20. Lee, C.; Beauchemin, K.A. A Review of Feeding Supplementary Nitrate to Ruminant Animals: Nitrate Toxicity, Methane Emissions, and Production Performance. *Can. J. Anim. Sci.* **2014**, *94*, 557–570. [[CrossRef](#)]
21. Pacheco, D.; Waghorn, G.; Janssen, P.H. Decreasing Methane Emissions from Ruminants Grazing Forages: A Fit with Productive and Financial Realities? *Anim. Prod. Sci.* **2014**, *54*, 1141–1154. [[CrossRef](#)]
22. Beauchemin, K.A.; McGinn, S.M. Methane Emissions from Feedlot Cattle Fed Barley or Corn Diets. *J. Anim. Sci.* **2005**, *83*, 653–661. [[CrossRef](#)]
23. Asanuma, N.; Iwamoto, M.; Hino, T. Effect of the Addition of Fumarate on Methane Production by Ruminal Microorganisms In Vitro. *J. Dairy Sci.* **1999**, *82*, 780–787. [[CrossRef](#)]
24. Tejido, M.; Ranilla, M.; García-Martínez, R.; Carro, M. In vitro microbial growth and rumen fermentation of different substrates as affected by the addition of disodium malate. *Anim. Sci.* **2005**, *81*, 31–38. [[CrossRef](#)]
25. Grainger, C.; Beauchemin, K.A. Can Enteric Methane Emissions from Ruminants Be Lowered without Lowering Their Production? *Anim. Feed Sci. Technol.* **2011**, *166–167*, 308–320. [[CrossRef](#)]
26. Carulla, J.E.; Kreuzer, M.; Machmüller, A.; Hess, H.D. Supplementation of Acacia Mearnsii Tannins Decreases Methanogenesis and Urinary Nitrogen in Forage-Fed Sheep. *Aust. J. Agric. Res.* **2005**, *56*, 961. [[CrossRef](#)]
27. Grainger, C.; Clarke, T.; Auldist, M.J.; Beauchemin, K.A.; McGinn, S.M.; Waghorn, G.C.; Eckard, R.J. Potential Use of Acacia Mearnsii Condensed Tannins to Reduce Methane Emissions and Nitrogen Excretion from Grazing Dairy Cows. *Can. J. Anim. Sci.* **2009**, *89*, 241–251. [[CrossRef](#)]
28. Benchaar, C.; Greathead, H. Essential Oils and Opportunities to Mitigate Enteric Methane Emissions from Ruminants. *Anim. Feed Sci. Technol.* **2011**, *166–167*, 338–355. [[CrossRef](#)]
29. McGinn, S.M.; Beauchemin, K.A.; Coates, T.; Colombatto, D. Methane Emissions from Beef Cattle: Effects of Monensin, Sunflower Oil, Enzymes, Yeast, and Fumaric Acid. *J. Anim. Sci.* **2004**, *82*, 3346–3356. [[CrossRef](#)] [[PubMed](#)]
30. McCaughey, W.P.; Wittenberg, K.; Corrigan, D. Impact of Pasture Type on Methane Production by Lactating Beef Cows. *Can. J. Anim. Sci.* **1999**, *79*, 221–226. [[CrossRef](#)]
31. Waghorn, G.C.; Tavendale, M.H.; Woodfield, D.R. Methanogenesis from Forages Fed to Sheep. In *Proceedings of the Conference New Zealand Grassland Association, Gore, NZ, USA, 12–16 November 2012*; pp. 167–172.
32. Allen, M.S. Physical Constraints on Voluntary Intake of Forages by Ruminants. *J. Anim. Sci.* **1996**, *74*, 3063. [[CrossRef](#)] [[PubMed](#)]
33. Meyer, K.; Hummel, J.; Clauss, M. The Relationship between Forage Cell Wall Content and Voluntary Food Intake in Mammalian Herbivores. *Mammal Rev.* **2010**, *40*, 221–245. [[CrossRef](#)]

34. Moss, A.R.; Jouany, J.-P.; Newbold, J. Methane Production by Ruminants: Its Contribution to Global Warming. *Ann. Zootech.* **2000**, *49*, 231–253. [CrossRef]
35. Ominski, K.H.; Wittenberg, K.M. Strategies for Reducing Enteric Methane Emissions in Forage-Based Beef Production Systems. In *Climate Change and Managed Ecosystems*; CRC Press: Boca Raton, FL, USA, 2005; pp. 261–272.
36. Pelletier, S.; Tremblay, G.F.; Bélanger, G.; Bertrand, A.; Castonguay, Y.; Pageau, D.; Drapeau, R. Forage Nonstructural Carbohydrates and Nutritive Value as Affected by Time of Cutting and Species. *Agron. J.* **2010**, *102*, 1388. [CrossRef]
37. Phelan, P.; Moloney, A.P.; McGeough, E.J.; Humphreys, J.; Bertilsson, J.; O’Riordan, E.G.; O’Kiely, P. Forage Legumes for Grazing and Conserving in Ruminant Production Systems. *Crit. Rev. Plant Sci.* **2015**, *34*, 281–326. [CrossRef]
38. Fulkerson, W.J.; Neal, J.S.; Clark, C.F.; Horadagoda, A.; Nandra, K.S.; Barchia, I. Nutritive Value of Forage Species Grown in the Warm Temperate Climate of Australia for Dairy Cows: Grasses and Legumes. *Livest. Sci.* **2007**, *107*, 253–264. [CrossRef]
39. Villalba, J.J.; Ates, S.; MacAdam, J.W. Non-Fiber Carbohydrates in Forages and Their Influence on Beef Production Systems. *Front. Sustain. Food Syst.* **2021**, *5*, 566338. [CrossRef]
40. Van Soest, P.J.V. *Nutritional Ecology of the Ruminant*; Cornell University Press: Ithaca, NY, USA, 2018; ISBN 978-1-5017-3235-5.
41. Waghorn, G.C.; Shelton, I.D.; Thomas, V.J. Particle Breakdown and Rumen Digestion of Fresh Ryegrass (*Lolium perenne* L.) and Lucerne (*Medicago sativa* L.) Fed to Cows during a Restricted Feeding Period. *Br. J. Nutr.* **1989**, *61*, 409–423. [CrossRef]
42. Sun, X.; Henderson, G.; Cox, F.; Molano, G.; Harrison, S.J.; Luo, D.; Janssen, P.H.; Pacheco, D. Lambs Fed Fresh Winter Forage Rape (*Brassica napus* L.) Emit Less Methane than Those Fed Perennial Ryegrass (*Lolium perenne* L.), and Possible Mechanisms behind the Difference. *PLoS ONE* **2015**, *10*, e0119697. [CrossRef] [PubMed]
43. Archimède, H.; Eugène, M.; Marie Magdeleine, C.; Boval, M.; Martin, C.; Morgavi, D.P.; Lecomte, P.; Doreau, M. Comparison of Methane Production between C3 and C4 Grasses and Legumes. *Anim. Feed Sci. Technol.* **2011**, *166–167*, 59–64. [CrossRef]
44. Pitcher, L.R. Beef Average Daily Gain and Enteric Methane Emissions on Birdsfoot Trefoil, Cicer Milkvech and Meadow Brome Pastures; All Graduate Theses and Dissertations 4015. Master’s Thesis, Utah State University, Logan, UT, USA, 2015. Available online: <https://digitalcommons.usu.edu/etd/401590> (accessed on 12 August 2021).
45. Berthiaume, R.; Benchaar, C.; Chaves, A.V.; Tremblay, G.F.; Castonguay, Y.; Bertrand, A.; Bélanger, G.; Michaud, R.; Lafrenière, C.; McAllister, T.A. Effects of Nonstructural Carbohydrate Concentration in Alfalfa on Fermentation and Microbial Protein Synthesis in Continuous Culture. *J. Dairy Sci.* **2010**, *93*, 693–700. [CrossRef]
46. Popp, J.D.; McCaughey, W.P.; Cohen, R.D.H.; McAllister, T.A.; Majak, W. Enhancing Pasture Productivity with Alfalfa: A Review. *Can. J. Plant Sci.* **2000**, *80*, 513–519. [CrossRef]
47. MacAdam, J.W.; Ward, R.E.; Griggs, T.C.; Min, B.R.; Aiken, G.E. Average Daily Gain and Blood Fatty Acid Composition of Cattle Grazing the Nonbloating Legumes Birdsfoot Trefoil and Cicer Milkvech in the Mountain West. *Prof. Anim. Sci.* **2011**, *27*, 574–583. [CrossRef]
48. MacAdam, J.; Villalba, J. Beneficial Effects of Temperate Forage Legumes That Contain Condensed Tannins. *Agriculture* **2015**, *5*, 475–491. [CrossRef]
49. Yost, M.; Allen, N.; Creech, E.; Putnam, D.; Gale, J.; Shewmaker, G. *Ten Reasons Why Alfalfa Is Highly Suitable for the West*; Utah State University Agriculture Extension; AG/Crops/2020-01pr.; Utah State University College of Agriculture and Applied Sciences: Logan, UT, USA, 2020; Available online: https://digitalcommons.usu.edu/cgi/viewcontent.cgi?article=3100&context=extension_curall (accessed on 12 August 2021).
50. Wang, Y.; Majak, W.; McAllister, T.A. Frothy Bloat in Ruminants: Cause, Occurrence, and Mitigation Strategies. *Anim. Feed Sci. Technol.* **2012**, *172*, 103–114. [CrossRef]
51. Majak, W.; McAllister, T.A.; McCartney, D.; Stanford, K.; Cheng, K.J. *Bloat in Cattle*; Alberta Agriculture Food and Rural Development, Information Packaging Center: Edmonton, AB, Canada, 2003; pp. 1–24.
52. Cameron, A.R.; Malmø, J. A Survey of the Efficacy of Sustained-release Monensin Capsules in the Control of Bloat in Dairy. *Cattle. Aust. Vet. J.* **1993**, *70*, 1–4. [CrossRef]
53. Thompson, D.J.; Brooke, B.M.; Garland, G.J.; Hall, J.W.; Majak, W. Effect of Stage of Growth of Alfalfa on the Incidence of Bloat in Cattle. *Can. J. Anim. Sci.* **2000**, *80*, 725–727. [CrossRef]
54. Julier, B.; Guines, F.; Emile, J.-C.; Huyghe, C. Variation in Protein Degradability in Dried Forage Legumes. *Anim. Res.* **2003**, *52*, 401–412. [CrossRef]
55. Abdoun, K.; Stumpff, F.; Martens, H. Ammonia and Urea Transport across the Rumen Epithelium: A Review. *Anim. Health Res. Rev.* **2006**, *7*, 43–59. [CrossRef] [PubMed]
56. Lobley, G.E.; Milano, G.D. Regulation of Hepatic Nitrogen Metabolism in Ruminants. *Proc. Nutr. Soc.* **1997**, *56*, 547–563. [CrossRef]
57. Calsamiglia, S.; Ferret, A.; Reynolds, C.K.; Kristensen, N.B.; van Vuuren, A.M. Strategies for Optimizing Nitrogen Use by Ruminants. *Animal* **2010**, *4*, 1184–1196. [CrossRef]
58. Provenza, F.D. Postingestive Feedback as an Elementary Determinant of Food Preference and Intake in Ruminants. *J. Range Manag.* **1995**, *48*, 2–17. [CrossRef]
59. Kohn, R.A.; Dinneen, M.M.; Russek-Cohen, E. Using Blood Urea Nitrogen to Predict Nitrogen Excretion and Efficiency of Nitrogen Utilization in Cattle, Sheep, Goats, Horses, Pigs, and Rats. *J. Anim. Sci.* **2005**, *83*, 879–889. [CrossRef]
60. Getachew, G.; Depeters, E.J.; Pittroff, W.; Putnam, D.H.; Dandekar, A.M. Review: Does Protein in Alfalfa Need Protection from Rumen Microbes? *Prof. Anim. Sci.* **2006**, *22*, 364–373. [CrossRef]
61. Dijkstra, J.; Oenema, O.; van Groenigen, J.W.; Spek, J.W.; van Vuuren, A.M.; Bannink, A. Diet Effects on Urine Composition of Cattle and N₂O Emissions. *Animal* **2013**, *7*, 292–302. [CrossRef] [PubMed]

62. Zonderland-Thomassen, M.A.; Liefvering, M.; Ledgard, S.F. Water Footprint of Beef Cattle and Sheep Produced in New Zealand: Water Scarcity and Eutrophication Impacts. *J. Clean. Prod.* **2014**, *73*, 253–262. [[CrossRef](#)]
63. Leip, A.; Billen, G.; Garnier, J.; Grizzetti, B.; Lassaletta, L.; Reis, S.; Simpson, D.; Sutton, M.A.; de Vries, W.; Weiss, F. Impacts of European Livestock Production: Nitrogen, Sulphur, Phosphorus and Greenhouse Gas Emissions, Land-Use, Water Eutrophication and Biodiversity. *Environ. Res. Lett.* **2015**, *10*, 115004. [[CrossRef](#)]
64. Oenema, O.; Wrage, N.; Velthof, G.L.; van Groenigen, J.W.; Dolfing, J.; Kuikman, P.J. Trends in Global Nitrous Oxide Emissions from Animal Production Systems. *Nutr. Cycl. Agroecosystems* **2005**, *72*, 51–65. [[CrossRef](#)]
65. Huang, T.; Gao, B.; Hu, X.-K.; Lu, X.; Well, R.; Christie, P.; Bakken, L.R.; Ju, X.-T. Ammonia-Oxidation as an Engine to Generate Nitrous Oxide in an Intensively Managed Calcareous Fluvo-Aquic Soil. *Sci. Rep.* **2015**, *4*, 3950. [[CrossRef](#)] [[PubMed](#)]
66. Pachauri, R.K.; Mayer, L. *Climate Change 2014: Synthesis Report*; Intergovernmental Panel on Climate Change, Ed.; Intergovernmental Panel on Climate Change: Geneva, Switzerland, 2015; ISBN 978-92-9169-143-2.
67. Bao, Y.; Zhou, K.; Zhao, G. Nitrous Oxide Emissions from the Urine of Beef Cattle as Regulated by Dietary Crude Protein and Gallic Acid. *J. Anim. Sci.* **2018**, *96*, 3699–3711. [[CrossRef](#)]
68. Cai, Y.; Chang, S.X.; Cheng, Y. Greenhouse Gas Emissions from Excreta Patches of Grazing Animals and Their Mitigation Strategies. *Earth-Sci. Rev.* **2017**, *171*, 44–57. [[CrossRef](#)]
69. Wang, Y.; McAllister, T.A.; Acharya, S. Condensed Tannins in Sainfoin: Composition, Concentration, and Effects on Nutritive and Feeding Value of Sainfoin Forage. *Crop Sci.* **2015**, *55*, 13. [[CrossRef](#)]
70. Lees, G.L.; Suttill, N.H.; Gruber, M.Y. Condensed Tannins in Sainfoin. 1. A Histological and Cytological Survey of Plant Tissues. *Can. J. Bot.* **1993**, *71*, 1147–1152. [[CrossRef](#)]
71. Sengul, S. Performance of Some Forage Grasses or Legumes and Their Mixtures under Dry Land Conditions. *Eur. J. Agron.* **2003**, *19*, 401–409. [[CrossRef](#)]
72. Karnezos, T.P.; Matches, A.G.; Brown, C.P. Spring Lamb Production on Alfalfa, Sainfoin, and Wheatgrass Pastures. *Agron. J.* **1994**, *86*, 497–502. [[CrossRef](#)]
73. Maughan, B.; Provenza, F.D.; Tansawat, R.; Maughan, C.; Martini, S.; Ward, R.; Clemensen, A.; Song, X.; Cornforth, D.; Villalba, J.J. Importance of Grass-Legume Choices on Cattle Grazing Behavior, Performance, and Meat Characteristics. *J. Anim. Sci.* **2014**, *92*, 2309–2324. [[CrossRef](#)] [[PubMed](#)]
74. Lagrange, S.; Beauchemin, K.A.; MacAdam, J.; Villalba, J.J. Grazing Diverse Combinations of Tanniferous and Non-Tanniferous Legumes: Implications for Beef Cattle Performance and Environmental Impact. *Sci. Total Environ.* **2020**, *746*, 140788. [[CrossRef](#)] [[PubMed](#)]
75. Lagrange, S.P. Influence of Forage Diversity and Condensed Tannins on Livestock Foraging Behavior, Production and Environmental Impact; All Graduate Theses and Dissertations 7813. Ph.D. Thesis, Utah State University, Logan, UT, USA, 2020. Available online: <https://digitalcommons.usu.edu/etd/7813/> (accessed on 12 August 2021).
76. Grabber, J.H.; Riday, H.; Cassida, K.A.; Griggs, T.C.; Min, D.H.; MacAdam, J.W. Yield, Morphological Characteristics, and Chemical Composition of European- and Mediterranean-Derived Birdsfoot Trefoil Cultivars Grown in the Colder Continental United States. *Crop Sci.* **2014**, *54*, 1893. [[CrossRef](#)]
77. Gibb, M.; Orr, R. Grazing Behaviour of Ruminants. *IGER Innov.* **1997**, *1*, 54–57.
78. Grabber, J.H.; Coblenz, W.K.; Riday, H.; Griggs, T.C.; Min, D.H.; MacAdam, J.W.; Cassida, K.A. Protein and Dry-Matter Degradability of European- and Mediterranean-Derived Birdsfoot Trefoil Cultivars Grown in the Colder Continental USA. *Crop Sci.* **2015**, *55*, 1356. [[CrossRef](#)]
79. MacAdam, J.W.; Griggs, T.C. *Irrigated Birdsfoot Trefoil Variety Trial: Forage Yield*; All Current Publications Paper 1337; Utah State University: Logan, UT, USA, 2013; Available online: https://digitalcommons.usu.edu/extension_curall/1337 (accessed on 12 August 2021).
80. Barry, T.N.; McNabb, W.C. The Implications of Condensed Tannins on the Nutritive Value of Temperate Forages Fed to Ruminants. *Br. J. Nutr.* **1999**, *81*, 263–272. [[CrossRef](#)]
81. Waghorn, G. Beneficial and Detrimental Effects of Dietary Condensed Tannins for Sustainable Sheep and Goat Production—Progress and Challenges. *Anim. Feed Sci. Technol.* **2008**, *147*, 116–139. [[CrossRef](#)]
82. Aboagye, I.A.; Beauchemin, K.A. Potential of Molecular Weight and Structure of Tannins to Reduce Methane Emissions from Ruminants: A Review. *Animals* **2019**, *9*, 856. [[CrossRef](#)]
83. Zeller, W.E. Activity, Purification, and Analysis of Condensed Tannins: Current State of Affairs and Future Endeavors. *Crop Sci.* **2019**, *59*, 886–904. [[CrossRef](#)]
84. Mueller-Harvey, I.; Bee, G.; Dohme-Meier, F.; Hoste, H.; Karonen, M.; Kölliker, R.; Lüscher, A.; Niderkorn, V.; Pellikaan, W.F.; Salminen, J.P.; et al. Benefits of Condensed Tannins in Forage Legumes Fed to Ruminants: Importance of Structure, Concentration, and Diet Composition. *Crop Sci.* **2019**, *59*, 861. [[CrossRef](#)]
85. Naumann, H.D.; Tedeschi, L.O.; Zeller, W.E.; Huntley, N.F. The Role of Condensed Tannins in Ruminant Animal Production: Advances, Limitations and Future Directions. *Rev. Bras. Zootec.* **2017**, *46*, 929–949. [[CrossRef](#)]
86. Lees, G.L.; Gruber, M.Y.; Suttill, N.H. Condensed Tannins in Sainfoin. II. Occurrence and Changes during Leaf Development. *Can. J. Bot.* **1995**, *73*, 1540–1547. [[CrossRef](#)]

87. Theodoridou, K.; Aufrère, J.; Andueza, D.; Pourrat, J.; Le Morvan, A.; Stringano, E.; Mueller-Harvey, I.; Baumont, R. Effects of Condensed Tannins in Fresh Sainfoin (*Onobrychis viciifolia*) on in Vivo and in Situ Digestion in Sheep. *Anim. Feed Sci. Technol.* **2010**, *160*, 23–38. [[CrossRef](#)]
88. Berard, N.C.; Wang, Y.; Wittenberg, K.M.; Krause, D.O.; Coulman, B.E.; McAllister, T.A.; Ominski, K.H. Condensed Tannin Concentrations Found in Vegetative and Mature Forage Legumes Grown in Western Canada. *Can. J. Plant Sci.* **2011**, *91*, 669–675. [[CrossRef](#)]
89. Aerts, R.J.; Barry, T.N.; McNabb, W.C. Polyphenols and Agriculture: Beneficial Effects of Proanthocyanidins in Forages. *Agric. Ecosyst. Environ.* **1999**, *75*, 1–12. [[CrossRef](#)]
90. Jonker, A.; Yu, P. The Occurrence, Biosynthesis, and Molecular Structure of Proanthocyanidins and Their Effects on Legume Forage Protein Precipitation, Digestion and Absorption in the Ruminant Digestive Tract. *Int. J. Mol. Sci.* **2017**, *18*, 1105. [[CrossRef](#)]
91. Min, B.R.; McNabb, W.C.; Barry, T.N.; Peters, J.S. Solubilization and Degradation of Ribulose-1,5-Bisphosphate Carboxylase/Oxygenase (EC 4.1.1.39; Rubisco) Protein from White Clover (*Trifolium repens*) and Lotus Corniculatus by Rumen Microorganisms and the Effect of Condensed Tannins on These Processes. *J. Agric. Sci.* **2000**, *134*, 305–317. [[CrossRef](#)]
92. Jones, G.A.; McAllister, T.A. Effects of Sainfoin (*Onobrychis viciifolia* Scop.) Condensed Tannins on Growth and Proteolysis by Four Strains of Ruminal Bacteria. *Appl. Environ. Microbiol.* **1994**, *60*, 5. [[CrossRef](#)] [[PubMed](#)]
93. McNabb, W.C.; Waghorn, G.C.; Peters, J.S.; Barry, T.N. The Effect of Condensed Tannins in Lotus Pedunculatus on the Solubilization and Degradation of Ribulose-1,5-Bisphosphate Carboxylase (EC 4.1.1.39; Rubisco) Protein in the Rumen and the Sites of Rubisco Digestion. *Br. J. Nutr.* **1996**, *76*, 535–549. [[CrossRef](#)] [[PubMed](#)]
94. Aufrère, J.; Dudilieu, M.; Andueza, D.; Poncet, C.; Baumont, R. Mixing Sainfoin and Lucerne to Improve the Feed Value of Legumes Fed to Sheep by the Effect of Condensed Tannins. *Animal* **2013**, *7*, 82–92. [[CrossRef](#)] [[PubMed](#)]
95. Avila, S.C.; Kozloski, G.V.; Orlandi, T.; Mezzomo, M.P.; Stefanello, S. Impact of a Tannin Extract on Digestibility, Ruminal Fermentation and Duodenal Flow of Amino Acids in Steers Fed Maize Silage and Concentrate Containing Soybean Meal or Canola Meal as Protein Source. *J. Agric. Sci.* **2015**, *153*, 943–953. [[CrossRef](#)]
96. Perez-Maldonado, R.A.; Norton, B.W.; Kerven, G.L. Factors Affecting in Vitro Formation of Tannin-protein Complexes. *J. Sci. Food Agric.* **1995**, *69*, 291–298. [[CrossRef](#)]
97. Bermingham, E.N.; Hutchinson, K.J.; Revell, D.K.; Brookes, I.M.; McNabb, W.C. The Effect of Condensed Tannins in Sainfoin (*Onobrychis viciifolia*) and Sulla (*Hedysarum coronarium*) on the Digestion of Amino Acids in Sheep. *Proc. N. Z. Soc. Anim. Prod.* **2001**, *61*, 5.
98. Naumann, H.D.; Hagerman, A.E.; Lambert, B.D.; Muir, J.P.; Tedeschi, L.O.; Kothmann, M.M. Molecular Weight and Protein-Precipitating Ability of Condensed Tannins from Warm-Season Perennial Legumes. *J. Plant Interact.* **2014**, *9*, 212–219. [[CrossRef](#)]
99. Ropiak, H.M.; Lachmann, P.; Ramsay, A.; Green, R.J.; Mueller-Harvey, I. Identification of Structural Features of Condensed Tannins That Affect Protein Aggregation. *PLoS ONE* **2017**, *12*, e0170768. [[CrossRef](#)]
100. Aufrère, J.; Theodoridou, K.; Mueller-Harvey, I.; Yu, P.; Andueza, D. Ruminal Dry Matter and Nitrogen Degradation in Relation to Condensed Tannin and Protein Molecular Structures in Sainfoin (*Onobrychis viciifolia*) and Lucerne (*Medicago sativa*). *J. Agric. Sci.* **2014**, *152*, 333–345. [[CrossRef](#)]
101. Le Bourvellec, C.; Renard, C.M.G.C. Interactions between Polyphenols and Macromolecules: Quantification Methods and Mechanisms. *Crit. Rev. Food Sci. Nutr.* **2012**, *52*, 213–248. [[CrossRef](#)]
102. McAllister, T.A.; Martinez, T.; Bae, H.D.; Muir, A.D.; Yanke, L.J.; Jones, G.A. Characterization of Condensed Tannins Purified from Legume Forages: Chromophore Production, Protein Precipitation, and Inhibitory Effects on Cellulose Digestion. *J. Chem. Ecol.* **2005**, *31*, 2049–2068. [[CrossRef](#)]
103. Aufrère, J.; Dudilieu, M.; Poncet, C. In Vivo and in Situ Measurements of the Digestive Characteristics of Sainfoin in Comparison with Lucerne Fed to Sheep as Fresh Forages at Two Growth Stages and as Hay. *Animal* **2008**, *2*, 1331–1339. [[CrossRef](#)] [[PubMed](#)]
104. Rufino-Moya, P.J.; Blanco, M.; Bertolín, J.R.; Joy, M. Methane Production of Fresh Sainfoin, with or without PEG, and Fresh Alfalfa at Different Stages of Maturity is Similar but the Fermentation End Products Vary. *Animals* **2019**, *9*, 197. [[CrossRef](#)]
105. Scharenberg, A.; Arrigo, Y.; Gutzwiller, A.; Wyss, U.; Hess, H.D.; Kreuzer, M.; Dohme, F. Effect of Feeding Dehydrated and Ensiled Tanniferous Sainfoin (*Onobrychis viciifolia*) on Nitrogen and Mineral Digestion and Metabolism of Lambs. *Arch. Anim. Nutr.* **2007**, *61*, 390–405. [[CrossRef](#)] [[PubMed](#)]
106. Theodoridou, K.; Aufrère, J.; Andueza, D.; Le Morvan, A.; Picard, F.; Pourrat, J.; Baumont, R. Effects of Condensed Tannins in Wrapped Silage Bales of Sainfoin (*Onobrychis viciifolia*) on in Vivo and in Situ Digestion in Sheep. *Animal* **2012**, *6*, 245–253. [[CrossRef](#)]
107. Silanikove, N.; Perevolotsky, A.; Provenza, F.D. Use of tannin-binding chemicals to assay for tannins and their negative post-ingestive effects in ruminants. *Anim. Feed Sci. Technol.* **2001**, *91*, 69–81. [[CrossRef](#)]
108. Williams, C.M.; Eun, J.-S.; MacAdam, J.W.; Young, A.J.; Fellner, V.; Min, B.R. Effects of Forage Legumes Containing Condensed Tannins on Methane and Ammonia Production in Continuous Cultures of Mixed Ruminal Microorganisms. *Anim. Feed Sci. Technol.* **2011**, *166–167*, 364–372. [[CrossRef](#)]
109. Grosse Brinkhaus, A.; Bee, G.; Silacci, P.; Kreuzer, M.; Dohme-Meier, F. Effect of Exchanging *Onobrychis viciifolia* and *Lotus corniculatus* for *Medicago sativa* on Ruminal Fermentation and Nitrogen Turnover in Dairy Cows. *J. Dairy Sci.* **2016**, *99*, 4384–4397. [[CrossRef](#)]

110. Aufrère, J.; Dudilieu, M.; Poncet, C.; Baumont, R. Effect of Condensed Tannins in Sainfoin on in Vitro Protein Solubility of Lucerne. In Proceedings of the XX International Grassland Congress: Grasslands—A Global Resource, Dublin, Ireland, 31 August 2005; O'Mara, F.P., Wilkins, R.J., Mannelje, L., Lovett, D.K., Rogers, P.A.M., Boland, T.M., Eds.; Wageningen Academic Publishers: Wageningen, The Netherlands, 2005; p. 248.
111. Stewart, E.K.; Beauchemin, K.A.; Dai, X.; MacAdam, J.W.; Christensen, R.G.; Villalba, J.J. Effect of tannin-containing hays on enteric methane emissions and nitrogen partitioning in beef cattle. *J. Anim. Sci.* **2019**, *97*, 3286–3299. [[CrossRef](#)]
112. McNabb, W.C.; Peters, J.S.; Foo, L.Y.; Waghorn, G.C.; Jackson, F.S. Effect of Condensed Tannins Prepared from Several Forages on the in Vitro Precipitation of Ribulose-1,5-Bisphosphate Carboxylase (Rubisco) Protein and its Digestion by Trypsin (EC 2.4.21.4) and Chymotrypsin (EC 2.4.21.1). *J. Sci. Food Agric.* **1998**, *77*, 201–212. [[CrossRef](#)]
113. Lagrange, S.; Villalba, J.J. Tannin-Containing Legumes and Forage Diversity Influence Foraging Behavior, Diet Digestibility, and Nitrogen Excretion by Lambs. *J. Anim. Sci.* **2019**, *97*, 3994–4009. [[CrossRef](#)] [[PubMed](#)]
114. Azuhnw, B.N.; Hertzberg, H.; Arrigo, Y.; Gutzwiller, A.; Hess, H.D.; Mueller-Harvey, I.; Torgerson, P.R.; Kreuzer, M.; Dohme-Meier, F. Investigation of Sainfoin (*Onobrychis viciifolia*) Cultivar Differences on Nitrogen Balance and Fecal Egg Count in Artificially Infected Lambs. *J. Anim. Sci.* **2013**, *91*, 2343–2354. [[CrossRef](#)] [[PubMed](#)]
115. Patra, A.K.; Saxena, J. A New Perspective on the Use of Plant Secondary Metabolites to Inhibit Methanogenesis in the Rumen. *Phytochemistry* **2010**, *71*, 1198–1222. [[CrossRef](#)] [[PubMed](#)]
116. Min, B.R.; McNABB, W.C.; Barry, T.N.; Kemp, P.D.; Waghorn, G.C.; McDonald, M.F. The Effect of Condensed Tannins in Lotus Corniculatus upon Reproductive Efficiency and Wool Production in Sheep during Late Summer and Autumn. *J. Agric. Sci.* **1999**, *132*, 323–334. [[CrossRef](#)]
117. Christensen, R.G.; Yang, S.Y.; Eun, J.-S.; Young, A.J.; Hall, J.O.; MacAdam, J.W. Effects of Feeding Birdsfoot Trefoil Hay on Neutral Detergent Fiber Digestion, Nitrogen Utilization Efficiency, and Lactational Performance by Dairy Cows. *J. Dairy Sci.* **2015**, *98*, 7982–7992. [[CrossRef](#)] [[PubMed](#)]
118. Hunt, S.R.; MacAdam, J.W.; Griggs, T.C. Lignification and Tannin Localization during the Development of Birdsfoot Trefoil Stems. *Crop Sci. Madison* **2014**, *54*, 1876–1886. [[CrossRef](#)]
119. Hunt, S.R.; Griggs, T.C.; MacAdam, J.W. Change in Birdsfoot Trefoil (*Lotus corniculatus* L.) Nutritive Value with Stem Elongation, Flowering and Pod Formation. In *EGF at 50: The Future of European Grasslands, Proceedings of the 25th General Meeting of the European Grassland Federation, Aberystwyth, Wales, 7–11 September 2014*; IBERS, Aberystwyth University: Aberystwyth, UK, 2014; pp. 884–886.
120. Douglas, G.B.; Wang, Y.; Waghorn, G.C.; Barry, T.N.; Purchas, R.W.; Foote, A.G.; Wilson, G.F. Liveweight Gain and Wool Production of Sheep Grazing Lotus Corniculatus and Lucerne (*Medicago sativa*). *N. Z. J. Agric. Res.* **1995**, *38*, 95–104. [[CrossRef](#)]
121. Harris, S.L.; Clark, D.A.; Laboyrie, P.J. *Birdsfoot Trefoil—An Alternative Legume for New Zealand Dairy Pastures*; New Zealand Grassland Association: Wellington, New Zealand, 1998; Volume 60, pp. 99–103.
122. Woodward, S.L.; Waghorn, G.C.; Laboyrie, P.G. Condensed Tannins in Birdsfoot Trefoil (*Lotus corniculatus*) Reduce Methane Emissions from Dairy Cows. *Proc. N. Z. Soc. Anim. Prod.* **2004**, *64*, 6.
123. Moreira, G.D.; Lima, P.D.M.T.; Borges, B.O.; Primavesi, O.; Longo, C.; McManus, C.; Abdalla, A.; Louvandini, H. Tropical Tanniferous Legumes Used as an Option to Mitigate Sheep Enteric Methane Emission. *Trop. Anim. Health Prod.* **2013**, *45*, 879–882. [[CrossRef](#)]
124. Wang, S.; Terranova, M.; Kreuzer, M.; Marquardt, S.; Eggerschwiler, L.; Schwarm, A. Supplementation of Pelleted Hazel (*Corylus avellana*) Leaves Decreases Methane and Urinary Nitrogen Emissions by Sheep at Unchanged Forage Intake. *Sci. Rep.* **2018**, *8*, 1–10.
125. Piñero-Vázquez, A.T.; Jiménez-Ferrer, G.; Alayon-Gamboa, J.A.; Chay-Canul, A.J.; Ayala-Burgos, A.J.; Aguilar-Pérez, C.F.; Ku-Vera, J.C. Effects of Quebracho Tannin Extract on Intake, Digestibility, Rumen Fermentation, and Methane Production in Crossbred Heifers Fed Low-Quality Tropical Grass. *Trop. Anim. Health Prod.* **2018**, *50*, 29–36. [[CrossRef](#)]
126. Jayanegara, A.; Goel, G.; Makkar, H.P.S.; Becker, K. Divergence between Purified Hydrolysable and Condensed Tannin Effects on Methane Emission, Rumen Fermentation and Microbial Population in Vitro. *Anim. Feed Sci. Technol.* **2015**, *209*, 60–68. [[CrossRef](#)]
127. Hatew, B.; Stringano, E.; Mueller-Harvey, I.; Hendriks, W.H.; Carbonero, C.H.; Smith, L.M.J.; Pellikaan, W.F. Impact of Variation in Structure of Condensed Tannins from Sainfoin (*Onobrychis viciifolia*) on in Vitro Ruminant Methane Production and Fermentation Characteristics. *J. Anim. Physiol. Anim. Nutr.* **2016**, *100*, 348–360. [[CrossRef](#)]
128. McMahon, L.R.; Majak, W.; McAllister, T.A.; Hall, J.W.; Jones, G.A.; Popp, J.D.; Cheng, K.J. Effect of Sainfoin on in Vitro Digestion of Fresh Alfalfa and Bloat in Steers. *Can. J. Anim. Sci.* **1999**, *79*, 203–212. [[CrossRef](#)]
129. Theodoridou, K.; Aufrère, J.; Niderkorn, V.; Andueza, D.; Le Morvan, A.; Picard, F.; Baumont, R. In Vitro Study of the Effects of Condensed Tannins in Sainfoin on the Digestive Process in the Rumen at Two Vegetation Cycles. *Anim. Feed Sci. Technol.* **2011**, *170*, 147–159. [[CrossRef](#)]
130. Niderkorn, V.; Barbier, E.; Macheboeuf, D.; Torrent, A.; Mueller-Harvey, I.; Hoste, H. In Vitro Rumen Fermentation of Diets with Different Types of Condensed Tannins Derived from Sainfoin (*Onobrychis viciifolia* Scop.) Pellets and Hazelnut (*Corylus avellana* L.) Pericarps. *Anim. Feed Sci. Technol.* **2020**, *259*, 114357. [[CrossRef](#)]
131. Tavendale, M.H.; Meagher, L.P.; Pacheco, D.; Walker, N.; Attwood, G.T.; Sivakumaran, S. Methane Production from in Vitro Rumen Incubations with Lotus Pedunculatus and Medicago Sativa, and Effects of Extractable Condensed Tannin Fractions on Methanogenesis. *Anim. Feed Sci. Technol.* **2005**, *123–124*, 403–419. [[CrossRef](#)]

132. Saminathan, M.; Sieo, C.C.; Gan, H.M.; Abdullah, N.; Wong, C.M.V.L.; Ho, Y.W. Effects of Condensed Tannin Fractions of Different Molecular Weights on Population and Diversity of Bovine Rumen Methanogenic Archaea in Vitro, as Determined by High-Throughput Sequencing. *Anim. Feed Sci. Technol.* **2016**, *216*, 146–160. [[CrossRef](#)]
133. Tan, H.Y.; Sieo, C.C.; Abdullah, N.; Liang, J.B.; Huang, X.D.; Ho, Y.W. Effects of Condensed Tannins from *Leucaena* on Methane Production, Rumen Fermentation and Populations of Methanogens and Protozoa in Vitro. *Anim. Feed Sci. Technol.* **2011**, *169*, 185–193. [[CrossRef](#)]
134. Bodas, R.; Prieto, N.; García-González, R.; Andrés, S.; Giráldez, F.J.; López, S. Manipulation of Rumen Fermentation and Methane Production with Plant Secondary Metabolites. *Anim. Feed Sci. Technol.* **2012**, *176*, 78–93. [[CrossRef](#)]
135. Vasta, V.; Daghighi, M.; Cappucci, A.; Buccioni, A.; Serra, A.; Viti, C.; Mele, M. Invited Review: Plant Polyphenols and Rumen Microbiota Responsible for Fatty Acid Biohydrogenation, Fiber Digestion, and Methane Emission: Experimental Evidence and Methodological Approaches. *J. Dairy Sci.* **2019**, *102*, 3781–3804. [[CrossRef](#)]
136. Bae, H.D.; McAllister, T.A.; Yanke, J.; Cheng, K.J.; Muir, A.D. Effects of Condensed Tannins on Endoglucanase Activity and Filter Paper Digestion by *Fibrobacter Succinogenes* S85t. *Appl. Environ. Microbiol.* **1993**, *59*, 2132–2138. [[CrossRef](#)]
137. McSweeney, C.S.; Palmer, B.; Bunch, R.; Krause, D.O. Effect of the Tropical Forage *Calliandra* on Microbial Protein Synthesis and Ecology in the Rumen. *J. Appl. Microbiol.* **2001**, *90*, 78–88. [[CrossRef](#)]
138. Bento, M.H.L.; Acamovic, T.; Makkar, H.P.S. The Influence of Tannin, Pectin and Polyethylene Glycol on Attachment of ¹⁵N-Labelled Rumen Microorganisms to Cellulose. *Anim. Feed Sci. Technol.* **2005**, *122*, 41–57. [[CrossRef](#)]
139. Chung, Y.-H.; McGeough, E.J.; Acharya, S.; McAllister, T.A.; McGinn, S.M.; Harstad, O.M.; Beauchemin, K.A. Enteric Methane Emission, Diet Digestibility, and Nitrogen Excretion from Beef Heifers Fed Sainfoin or Alfalfa. *J. Anim. Sci.* **2013**, *91*, 4861–4874. [[CrossRef](#)]
140. Vaithiyanathan, S.; Bhatta, R.; Mishra, A.S.; Prasad, R.; Verma, D.L.; Singh, N.P. Effect of Feeding Graded Levels of *Prosopis Cineraria* Leaves on Rumen Ciliate Protozoa, Nitrogen Balance and Microbial Protein Supply in Lambs and Kids. *Anim. Feed Sci. Technol.* **2007**, *133*, 177–191. [[CrossRef](#)]
141. Bhatta, R.; Uyeno, Y.; Tajima, K.; Takenaka, A.; Yabumoto, Y.; Nonaka, I.; Enishi, O.; Kurihara, M. Difference in the Nature of Tannins on in Vitro Ruminant Methane and Volatile Fatty Acid Production and on Methanogenic Archaea and Protozoal Populations. *J. Dairy Sci.* **2009**, *92*, 5512–5522. [[CrossRef](#)] [[PubMed](#)]
142. Min, B.R.; Parker, D.; Brauer, D.; Waldrip, H.; Lockard, C.; Hales, K.; Akbay, A.; Augyte, S. The role of seaweed as a potential dietary supplementation for enteric methane mitigation in ruminants: Challenges and opportunities. *Anim. Nutr.* **2021**, *7*, 1371–1387. [[CrossRef](#)]
143. Ku-Vera, J.C.; Jiménez-Ocampo, R.; Valencia-Salazar, S.S.; Montoya-Flores, M.D.; Molina-Botero, I.C.; Arango, J.; Gómez-Bravo, C.A.; Aguilar-Pérez, C.F.; Solorio-Sánchez, F.J. Role of secondary plant metabolites on enteric methane mitigation in ruminants. *Front. Vet. Sci.* **2020**, *7*, 584. [[CrossRef](#)] [[PubMed](#)]
144. Mbiriri, D.T.; Cho, S.; Mamvura, C.I.; Choi, N.J. Assessment of rumen microbial adaptation to garlic oil, carvacrol and thymol using the consecutive batch culture system. *J. Vet. Sci. Anim. Husb.* **2015**, *4*, 1–7. [[CrossRef](#)]
145. McMahan, L.R.; McAllister, T.A.; Berg, B.P.; Majak, W.; Acharya, S.N.; Popp, J.D.; Coulman, B.E.; Wang, Y.; Cheng, K.J. A Review of the Effects of Forage Condensed Tannins on Ruminant Fermentation and Bloat in Grazing Cattle. *Can. J. Plant Sci.* **2000**, *80*, 469–485. [[CrossRef](#)]
146. Li, Y.-G.; Tanner, G.; Larkin, P. The DMACA–HCl Protocol and the Threshold Proanthocyanidin Content for Bloat Safety in Forage Legumes. *J. Sci. Food Agric.* **1996**, *70*, 89–101. [[CrossRef](#)]
147. Wang, Y.; Berg, B.P.; Barbieri, L.R.; Veira, D.M.; McAllister, T.A. Comparison of Alfalfa and Mixed Alfalfa-Sainfoin Pastures for Grazing Cattle: Effects on Incidence of Bloat, Ruminant Fermentation, and Feed Intake. *Can. J. Anim. Sci.* **2006**, *86*, 383–392. [[CrossRef](#)]
148. Chail, A.; Legako, J.F.; Pitcher, L.R.; Griggs, T.C.; Ward, R.E.; Martini, S.; MacAdam, J.W. Legume Finishing Provides Beef with Positive Human Dietary Fatty Acid Ratios and Consumer Preference Comparable with Grain-Finished Beef. *J. Anim. Sci.* **2016**, *94*, 2184–2197. [[CrossRef](#)]
149. Simopoulos, A.P. Omega-3 Fatty Acids in Inflammation and Autoimmune Diseases. *J. Am. Coll. Nutr.* **2002**, *21*, 495–505. [[CrossRef](#)] [[PubMed](#)]
150. Wall, R.; Ross, R.P.; Fitzgerald, G.F.; Stanton, C. Fatty Acids from Fish: The Anti-Inflammatory Potential of Long-Chain Omega-3 Fatty Acids. *Nutr. Rev.* **2010**, *68*, 280–289. [[CrossRef](#)]
151. Gomez Candela, C.; López, L.B.; Kohen, V.L. Importance of a Balanced Omega 6/Omega 3 Ratio for the Maintenance of Health Nutritional Recommendations. *Nutr. Hosp.* **2011**, *26*, 323–329. [[CrossRef](#)] [[PubMed](#)]
152. Vasta, V.; Makkar, H.P.S.; Mele, M.; Priolo, A. Ruminant Biohydrogenation as Affected by Tannins in Vitro. *Br. J. Nutr.* **2008**, *102*, 82–92. [[CrossRef](#)] [[PubMed](#)]
153. Vasta, V.; Mele, M.; Serra, A.; Scerra, M.; Luciano, G.; Lanza, M.; Priolo, A. Metabolic Fate of Fatty Acids Involved in Ruminant Biohydrogenation in Sheep Fed Concentrate or Forage with or without Tannins. *J. Anim. Sci.* **2009**, *87*, 2674–2684. [[CrossRef](#)]
154. Temperton, V.M.; Mwangi, P.N.; Scherer-Lorenzen, M.; Schmid, B.; Buchmann, N. Positive interactions between nitrogen-fixing legumes and four different neighboring species in a biodiversity experiment. *Oecologia* **2007**, *151*, 190–205. [[CrossRef](#)]
155. Pirhofer-Walzl, K.; Rasmussen, J.; Høgh-Jensen, H.; Eriksen, J.; Søgaard, K.; Rasmussen, J. Nitrogen transfer from forage legumes to nine neighbouring plants in a multi-species grassland. *Plant Soil* **2012**, *350*, 71–84. [[CrossRef](#)]

156. Rochette, P.; Janzen, H.H. Towards a revised coefficient for estimating N₂O emissions from legumes. *Nutr. Cycl. Agroecosyst.* **2005**, *73*, 171–179. [CrossRef]
157. Tilman, D. *Resource Competition and Community Structure*; Princeton University Press: Princeton, NJ, USA, 1982; ISBN 0-691-08302-9.
158. Provenza, F.D.; Villalba, J.J.; Haskell, J.; MacAdam, J.W.; Griggs, T.C.; Wiedmeier, R.D. The Value to Herbivores of Plant Physical and Chemical Diversity in Time and Space. *Crop Sci.* **2007**, *47*, 382. [CrossRef]
159. Westoby, M. What Are the Biological Bases of Varied Diets? *Am. Nat.* **1978**, *112*, 627–631. [CrossRef]
160. Villalba, J.J.; Provenza, F.D.; Catanese, F.; Distel, R.A. Understanding and Manipulating Diet Choice in Grazing Animals. *Anim. Prod. Sci.* **2015**, *55*, 261. [CrossRef]
161. Waghorn, G.C.; McNabb, W.C. Consequences of Plant Phenolic Compounds for Productivity and Health of Ruminants. *Proc. Nutr. Soc.* **2003**, *62*, 383–392. [CrossRef] [PubMed]
162. Rochfort, S.; Parker, A.J.; Dunshea, F.R. Plant Bioactives for Ruminant Health and Productivity. *Phytochemistry* **2008**, *69*, 299–322. [CrossRef] [PubMed]
163. Niderkorn, V.; Mueller-Harvey, I.; Le Morvan, A.; Aufrère, J. Synergistic Effects of Mixing Cocksfoot and Sainfoin on In Vitro Rumen Fermentation. Role of Condensed Tannins. *Anim. Feed Sci. Technol.* **2012**, *178*, 48–56. [CrossRef]
164. Sinz, S.; Marquardt, S.; Soliva, C.R.; Braun, U.; Liesegang, A.; Kreuzer, M. Phenolic Plant Extracts are Additive in their Effects Against in Vitro Ruminal Methane and Ammonia Formation. *Asian-Australas. J. Anim. Sci.* **2019**, *32*, 966–976. [CrossRef]
165. Aufrère, J.; Dudillieu, M.; Poncet, C.; Baumont, R.; Dumont, B. *Effect of Condensed Tannins in Sainfoin on In Vitro Protein Solubility of Lucerne as Affected by the Proportion of Sainfoin in the Mixture and the Preserving Conditions*; Serie A; Options Méditerranéennes: Paris, France, 2007; pp. 63–66. Available online: <http://om.ciheam.org/article.php?IDPDF=800355> (accessed on 12 August 2021).
166. Villalba, J.J.; Provenza, F.D.; Han, G. Experience Influences Diet Mixing by Herbivores: Implications for Plant Biochemical Diversity. *Oikos* **2004**, *107*, 100–109. [CrossRef]
167. Rogosic, J.; Estell, R.E.; Skobic, D.; Stanic, S. Influence of Secondary Compound Complementarity and Species Diversity on Consumption of Mediterranean Shrubs by Sheep. *Appl. Anim. Behav. Sci.* **2007**, *107*, 58–65. [CrossRef]
168. Meuret, M.; Bruchou, C. Modélisation de l'ingestion Selon La Diversité Des Choix Alimentaires Réalisés Par La Chèvre Au Pâturage Sur Parcours. *Rencontres Rech. Rumin.* **1994**, *1*, 225–228.
169. Provenza, F.D. Acquired Aversions as the Basis for Varied Diets of Ruminants Foraging on Rangelands. *J. Anim. Sci.* **1996**, *74*, 2010–2020. [CrossRef] [PubMed]
170. Christensen, R.G. Improvement of Nutrient Utilization Efficiency, Ruminal Fermentation and Lactational Performance of Dairy Cows by Feeding Birdsfoot Trefoil. Ph.D. Thesis, Utah State University, Logan, UT, USA, 2015. Available online: <https://digitalcommons.usu.edu/etd/4286207> (accessed on 15 August 2021).
171. Senft, R.L.; Coughenour, M.B.; Bailey, D.W.; Rittenhouse, L.R.; Sala, O.E.; Swift, D.M. Large Herbivore Foraging and Ecological Hierarchies. *BioScience* **1987**, *37*, 789–799. [CrossRef]
172. Chapman, D.F.; Parsons, A.J.; Cosgrove, G.P.; Barker, D.J.; Marotti, D.M.; Venning, K.J.; Rutter, S.M.; Hill, J.; Thompson, A.N. Impacts of Spatial Patterns in Pasture on Animal Grazing Behavior, Intake, and Performance. *Crop Sci.* **2007**, *47*, 399. [CrossRef]
173. Prache, S.J.; Gordon, I.; Rook, A.J. Foraging Behaviour and Diet Selection in Domestic Herbivores. *Ann. Zootech.* **1998**, *47*, 335–345. [CrossRef]
174. Acharya, S.; Sottie, E.; Coulman, B.; Iwaasa, A.; McAllister, T.; Wang, Y.; Liu, J. New Sainfoin Populations for Bloat-Free Alfalfa Pasture Mixtures in Western Canada. *Crop Sci.* **2013**, *53*, 2283–2293. [CrossRef]
175. Sottie, E.T.; Acharya, S.N.; McAllister, T.; Thomas, J.; Wang, Y.; Iwaasa, A. Alfalfa Pasture Bloat Can Be Eliminated by Intermixing with Newly-Developed Sainfoin Population. *Agron. J.* **2014**, *106*, 1470. [CrossRef]
176. Villalba, J.J.; Manteca, X. A Case for Eustress in Grazing Animals. *Front. Vet. Sci.* **2019**, *6*, 303. [CrossRef] [PubMed]
177. Catanese, F.; Obelar, M.; Villalba, J.J.; Distel, R.A. The Importance of Diet Choice on Stress-Related Responses by Lambs. *Appl. Anim. Behav. Sci.* **2013**, *148*, 37–45. [CrossRef]
178. Lyons, D.M.; Parker, K.J. Stress Inoculation-Induced Indications of Resilience in Monkeys. *J. Trauma. Stress.* **2007**, *20*, 423–433. [CrossRef]
179. Manteca, X.; Villalba, J.J.; Atwood, S.B.; Dziba, L.; Provenza, F.D. Is Dietary Choice Important to Animal Welfare? *J. Vet. Behav.* **2008**, *3*, 229–239. [CrossRef]
180. Sanderson, M.A.; Goslee, S.C.; Soder, K.J.; Skinner, R.H.; Tracy, B.F.; Deak, A. Plant Species Diversity, Ecosystem Function, and Pasture Management—A Perspective. *Can. J. Plant Sci.* **2007**, *87*, 479–487. [CrossRef]
181. Villalba, J.J.; Beauchemin, K.A.; Gregorini, P.; MacAdam, J.W. Pasture Chemoscapes and Their Ecological Services. *Transl. Anim. Sci.* **2019**, *3*, txz003. [CrossRef] [PubMed]
182. Caroprese, M.; Ciliberti, M.G.; Marino, R.; Napolitano, F.; Braghieri, A.; Sevi, A.; Albenzio, M. Effect of Information on Geographical Origin, Duration of Transport and Welfare Condition on Consumer's Acceptance of Lamb Meat. *Sci. Rep.* **2020**, *10*, 9754. [CrossRef] [PubMed]
183. Greenwood, P.L. Review: An Overview of Beef Production from Pasture and Feedlot Globally, as Demand for Beef and the Need for Sustainable Practices Increase. *Animal* **2021**, 100295. [CrossRef] [PubMed]