

PEANUT SCIENCE

The Journal of the American Peanut Research and Education Society

2022 Bailey Award Winner

ARTICLE

Discovery of a resistance gene cluster associated with smut resistance in peanut

K.D. Chamberlin¹; R.S. Bennett¹; J. Baldessari²; G. De La Barrera²; G. Cordes²; N.G. Grandon²; E.M.C. Mamani²; A.V. Rodriguez²; S. Morichetti³; C.C. Holbrook⁴; P. Ozias-Akins⁵; Y. Chu⁵; S.P. Tallury⁶; J. Clevenger⁷; W. Korani⁷; B. Scheffler⁸; R. Youngblood⁸; S. Simpson⁸

¹USDA-ARS, Peanut and Small Grains Research Unit, Stillwater, Oklahoma

²Instituto Nacional de Tecnologia Agropecuaria, Manfredi, Argentina

³Aceitera, General Deheza, Argentina

⁴USDA-ARS, Crop Genetics and Breeding Research Unit, Tifton, Georgia

⁵National Environmentally Sound Production Agriculture Laboratory, Institute of Plant Breeding, Genetics, and Genomics, University of Georgia, Tifton, Georgia

⁶USDA-ARS, Plant Genetic Resources Conservation Unit, Griffin, Georgia

⁷HudsonAlpha Institute for Biotechnology, Huntsville, Alabama

⁸USDA-ARS, Genomics and Bioinformatics Research Unit, Stoneville, Mississippi

ARTICLE INFORMATION

Keywords:

Peanut, Smut, Resistance, Quantitative Trait Loci

Corresponding Author email:

Kelly.Chamberlin@usda.gov

DOI: 10.3146/0095-3679-51-PS23-6

ABSTRACT

Peanut smut, caused by *Thecaphora frezzii* Carranza & J.C. Lindq., is an emerging threat for the global peanut industry. The disease's destructive potential can be exemplified by pod incidence values as high as 70% and yield losses reaching 30%. Because fungicides have shown moderate but highly variable levels of control levels, development and deployment of smut resistant cultivars are the best strategies for disease management. Screening for smut-resistant germplasm requires years of field trials and is currently the only option for breeders because genetic markers for resistance have not yet been developed. The objectives of this study were to perform whole genome sequencing (WGS) on populations developed for smut resistance mapping and subsequently fine map discovered QTL associated with smut resistance. An expedited strategy was employed by phenotyping in the F₃ generation. We phenotyped 200 families with 3 plants per family for smut resistance in infested fields during the 2019/2020 season in General Deheza (32°45'23"S 63°47'20"W), Argentina. Each individual was sequenced using iGenomXRipTide library preparation and Illumina NovaSeq sequencing to yield approximately 1 times genome coverage. Analysis of the phenotype and genotype data using Khufu resulted in the identification of a single major smut resistance QTL on chromosome 12 (B02). Chromosome level genome sequences were assembled for the resistant parent (Ascasubi) and susceptible parent (Granoleico) using PacBio HIFI sequencing. A validation population was sequenced using Khufu to validate the QTL region and analysis confirmed the major locus on chromosome 12. The identified variation will be used to develop smut resistant varieties quickly using molecular assisted breeding strategies.

INTRODUCTION

Peanut smut, caused by *Thecaphora frezzii* Carranza & J.C. Lindq. (*T. frezzii*), is an emerging threat for the global peanut industry (BICON, 2017; Rago *et al.*, 2017). First described in 1962, *T. frezzii* was observed on a collected sample of a wild peanut from Mato Grosso do Sul, Brazil, that was sent to INTA's Manfredi Exp. Stn. in Córdoba Province, Argentina (Carranza & Lindquist 1962). In 1995, peanut smut was first detected on cultivated peanuts on three farms in north-central Córdoba (Marinelli *et al.*, 2008). Since then, it has spread to all peanut-producing provinces in Argentina (Cazón *et al.*, 2018). The disease's destructive potential can be exemplified by pod incidence values as high as 70% (Bonessi *et al.* 2011, Chamberlin *et al.* 2022) and yield losses reaching 30% (Oddino *et al.* 2010). Disease management strategies using pesticides or biocontrol agents (Figueredo *et al.* 2017), liming (Bonessi *et al.* 2011), tillage (Cignetti *et al.* 2010; Cazón *et al.* 2014), and rotation (Marraro Acuña and Haro, 2011) have been attempted with modest results. Fungicides have shown moderate but highly variable control levels (Rago *et al.* 2017; Paredes *et al.* 2021). Smut resistant cultivars seem the best tool for disease management as they suffer less crop loss and limit the pathogen's multiplication (Rago *et al.* 2017). Sources of smut resistance have been reported in wild peanuts (de Blas *et al.* 2019), synthetic amphidiploids (de Blas *et al.* 2021) and in *A. hypogaea* core collections (Chamberlin *et al.* 2022; Wann *et al.* 2020). Smut genetic resistance has shown high broad sense heritability (Bressano *et al.* 2019; de Blas *et al.* 2021). Recently, major and a minor quantitative trait loci (QTL) associated with moderate resistance have been identified in a recombinant inbred line (RIL) population developed from a cross between a susceptible *A. hypogaea* genotype and a resistant synthetic amphidiploid [*A. correntina* × *A. cardenasii*] × *A. batizocoi* (de Blas *et al.* 2021). Peanut reaction to *T. frezzii* is assessed by opening each pod by hand and inspecting it visually for incidence (i.e., the presence or absence of *T. frezzii* in pods), which is time and labor intensive. Screening for smut resistant genetic sources requires years of in-field trials along with pod phenotyping and is currently the only option for breeders because markers for resistance have not yet been developed.

Marker assisted selection (MAS) is now widely used in plant breeding to increase the efficiency of developing cultivars with highly desired traits. Early marker work identified several for disease resistance in peanut. Random amplified polymorphic DNA (RAPD) was used to identify root-knot nematode (RKN) resistance (Burow *et al.*, 1996), restriction fragment length polymorphisms (RFLPs) were identified for resistance to *Meloidogyne arenaria* (Church *et al.*, 2000), amplified fragment length polymorphisms (AFLPs) were found linked to resistance to the vector of groundnut rosette virus (Herselman *et al.*, 2004), and single sequence repeats (SSRs) associated with resistance to *Sclerotinia minor* (Jagger) (Chenault, *et al.*, 2009) along with those for tomato spotted wilt virus (TSWV) and leaf spot (LS) (Wang *et al.*, 2013; Tseng *et al.*, 2016; Zhao *et al.*, 2018), rust (Khedikar *et al.*, 2010; Mondal and Badigannavar, 2018), and bud necrosis disease (Jadhav *et al.*, 2019) were identified.

Sequencing of the diploid progenitors (Bertioli *et al.*, 2016), *A. duranensis* and *A. ipaensis*, provided the reference

genomes necessary to construct the high-density single nucleotide polymorphism (SNP) genotyping array Axiom_Arachis (Pandey *et al.*, 2017) which led to the generation of SNP-based linkage maps enabling the identification SNPs associated with traits. SNPs are preferred over other marker types due to their wide distribution throughout the peanut genome (Liao and Lee, 2010). Whole genome resequencing (WGRS) of mapping populations has also been used to generate high-density SNP-based maps and identify QTL along with candidate genes for resistance to peanut diseases such as early LS (ELS), late LS (LLS), and spotted wilt (Agarwal *et al.*, 2018). Such linkage maps have also been used to identify QTL for resistance in peanut to RKN (Leal-Bertioli *et al.*, 2016), *Aspergillus flavus* (Khan *et al.*, 2020), bacterial wilt (Luo *et al.*, 2020a), stem rot (Luo *et al.* 2020b), Sclerotinia blight (Liang *et al.*, 2021), and bud necrosis disease (Jasani *et al.*, 2021). The recent release of the tetraploid (cultivated) peanut genome sequence (Bertioli *et al.*, 2019) has further clarified questions regarding *A. hypogaea* evolution and the generation of a high-density SNP array for tetraploid peanut (Axiom_Arachis2) (Clevenger *et al.*, 2017, Clevenger *et al.*, 2018) provide further foundation for rapid marker-trait association. Using the Axiom_Arachis SNP array, potential SNPs and QTL associated with peanut smut resistance were identified by association mapping using a limited number of genotypes (Massa *et al.*, 2021), but no confirmed QTL or candidate genes have been reported to date. Therefore, the objectives of this study were to perform whole genome sequencing (WGS) on a population developed for smut resistance mapping and subsequently fine map discovered QTL associated with smut resistance.

MATERIALS AND METHODS

Plant Materials and Population Development.

The mapping population consisting of 200 lines was developed by crossing Mf10_2870, smut susceptible (J. Baldessari, unpublished data), and Ascasubi Hispano, smut resistant (Ibañez *et al.*, 2018). An additional population used for validation purposes was also developed by crossing Granoleico, smut susceptible (Soave, 2002; Oddino *et al.*, 2013) and Ascasubi Hispano. To rapidly identify QTL associated with smut resistance, a non-traditional approach was used where families within the population were phenotyped and genotyped at familial stages prior to the F₇ generation rather than developing a uniform recombinant inbred line (RIL) population. Briefly, three individuals from each F_{2,3} family (for the mapping population) and from each F_{3,4} family (for the validation population) were phenotyped under heavily smut infested field conditions. This was done during the 2019-2020 season for the mapping population and during the 2020-2021 season for the validation population, in General Deheza, Córdoba Province, Argentina under rainfed conditions.

Experimental Design and Field Testing.

Smut resistance tests were conducted as described in Chamberlin *et al.*, 2022, in fields adjacent to a farm (-32.759, -63.770) in the town of General Deheza (Córdoba Province, Argentina). Soil at the site was a General Deheza coarse-silty sandy loam type (0.3% slope, coarse-silty, mixed, thermic, Entic Haplustoll). The tests were planted one km downwind from a

peanut processing plant where peanut smut is prevalent. The production field site was on a rotation schedule of corn-corn-soybean-peanut, where the preceding crop of the test site was always soybean.

Two control entries, one resistant (Ascasubi Hispano) and one susceptible (Colorado Irradiado INTA) were planted in both seasons. They were replicated twelve times each. Each family was represented by three individuals, each one constituting a plot. Plots were arranged in a square grid and randomized within it. Controls were placed, alternately, along a diagonal of the grid. Individual plots were spaced one m apart among rows and between plants.

The tests were planted on 11/7/19 and 12/23/20, using an augmented grid design with three replications, using single plants as experimental units. Plants were respectively dug at 147 and 140 days after planting (DAP), and all pods from each plant were harvested by hand and placed in mesh bags to air dry for two months. Yearly soil spore count was done as described by Chamberlin *et al.* (2022) to determine disease pressure. The test was conducted following the extension guidelines of the Instituto Nacional de Tecnología Agropecuaria (INTA).

Harvest and Evaluation of Smut Resistance.

Each pod was opened by hand and rated for disease incidence, i.e., presence or absence of *T. frezzii* sori or spore masses on the kernels. Disease incidence (DI) was calculated as DI = Infected pods / Total pods.

QTL mapping and validation.

DNA was taken from 572 individuals (approximately 3 individuals from each of the 200 lines) of the F_{2:3} family (Mf10_2870 and Ascasubi Hispano. DNA was also taken from 96 individual plants of F_{2:3} family of Ascasubi x Granoleico selected for phenotype. DNA extraction was performed from 20 mg of dried young leaves in silica gel. The samples were ground for two 10-sec cycles in bead mill at 15,000 rpm (Super FastPrep-2 Bead Beating System, MP BiomedicalsLLC, Irvine, CA, USA). A modified CTAB method was used with a sorbitol cleaning wash before the lysis step (Inglis *et al.* 2018). DNA quality was determined using 0.8% agarose gel and the DNA quantity was estimated with DS-11+ Nano UV-Vis Spectrophotometer (DeNovix Inc., DE, USA).

Parental genomic DNA extraction.

For whole-genome sequencing, seeds of Ascasubi INTA Cv, Granoleico Cv and Mf10_2870L were germinated in an incubation oven, under total darkness at constant temperature of 28 °C for 15 days. Fresh young etiolated leaves were collected for each genotype, immediately frozen with liquid nitrogen, and preserved at -80 C (Revco, Model ULT1386-5V41). The nuclei were extracted following the Nucleii Isolation – LN2 Plant Tissue Protocol (Ciculomics Inc./ PacBio) optimized from Workman *et al.* (2018). In the step 5 of protocol, filtration of the lysate with steriflip 20 µm pore was replaced with two lysate filtrations with a cell strainer of 100 µm pore and 40 µm pore. High-molecular weight (HMW) DNA was extracted from nuclei following the specifications of Nanobind Plant Nucleii Big DNA Kit (Ciculomics Inc./ PacBio). DNA quality was determined using 0.8% agarose gel and the DNA quantity was

estimated by Qubit V1 (Invitrogen ThermoFisher Scientific Inc.) using Qubit® dsDNA BR Assay Kits.

Genotyping: Affymetrix Axiom SNP Array.

Seeds from the peanut mini core accessions were sowed in 4.7-m pots with a mix of 50% Promix (Premier Tech Horticulture, Quaker, PA) and 50% steam-sterilized sandy soil from the Coastal Plain Experiment Station in Tifton, GA. Young leaf tissue was collected from 1-month-old plants for DNA extraction using DNeasy Plant mini kit (www.qiagen.com). Quantification of DNA was performed with Quant-iT dsDNA assay kit (www.thermofisher.com). DNA were submitted for genotyping with the *Arachis* version SNP array consisting of 47K SNPs features (www.thermofisher.com). SNP data were curated from the Axiom analysis suite (www.thermofisher.com). SNP markers were classified into six categories, i.e., PolyHighResolution, NoMinorHom, MonoHighResolution, CallRateBelowThreshold, OfftargetVariant, and Other according to the SNP QC matrix of the software (Clevenger *et al.*, 2017). All data analysis in this study was performed with markers in the PolyHighResolution category due to its high quality in signal separation.

Genotyping: PacBio Sequencing

High molecular weight DNA was received from *A. hypogea* Granoleico and *A. hypogea* Ascasubi. DNA was sheared using the Diagenode Megaruptor 3 targeting 20kb fragments. Sheared DNA was prepared for PacBio sequencing using the SMRTbell Express Template Prep Kit 2.0. The library was size selected with AMPure PB beads to remove fragments less than 3kb. Sequencing was performed on a Sequel IIe System (Pacific Biosciences, Menlo Park, CA) using Binding Kit 2.2, Sequel II Sequencing Kit 2.0, and SMRTCell 8M. To target HiFi reads, the library was sequenced using a 30-hour movie time using Instrument Control Software Version 10. Raw subreads were converted to HiFi data by processing with CCS to call a single high quality consensus sequence for each molecule, using a 99.5% consensus accuracy cutoff. *A. hypogea* Granoleico was sequenced on 2 SMRT cells yielding 73.11Gb of HiFi data, and *A. hypogea* Ascasubi was sequenced on 3 SMRT cells yielding 61.37Gb of HiFi data.

Genotyping: Khufu whole-genome sequencing and analysis

DNA from populations were prepped into sequencing libraries using iGenomX Riptide library prep. For the F_{2:3} population, Six Riptide libraries, representing 576 samples were sequenced on 2 lanes of NovaSeq S4 chemistry (800 Gb raw base pairs per lane) to yield an average of 0.99X genome coverage per individual. The raw reads were demultiplexed and processed using Khufu (hudsonalpha.org/khufudata). To identify QTL for resistance, bulks were identified using one year of field data. Analysis of the bulks were done *in silico*, by bulking individual sequenced samples and calculating allele frequency differences between the putative resistant and susceptible bulks. The F_{3:4} samples were processed and analyzed the same way, except were sequenced to an average of 1.14X genome coverage. Haplotype analysis was done using HawkHAP, which is a component of Khufu.

RESULTS AND DISCUSSION

Generation of reference genomes

Long read sequencing and assembly led to the generation of highly contiguous genome assemblies for smut resistant parent Ascasubi, and smut susceptible parent Granoleico (Table 1).

Mapping of smut resistance

A total of 576 individuals to construct bulks for mapping. They represented 200 F2:3 sub-families where each family was represented by 3 individuals. The smut pressure was low due to drought stress during pod development, but there were enough 'susceptible' individuals to map construct bulks (Figure 1). The frequency of infection was heavily skewed to the low end of the distribution, indicating that there was a considerable amount of false negative disease scores. To construct bulks, we employed a conservative approach whereby we selected susceptible individuals when all three sub-family members had infection and the average infection was greater than 5%. It was more difficult to select resistant bulks, because more than 400 individuals had 0% infection. For the 'resistant' bulk we selected every family member if they each had 0% infection and at least one individual was located in the field close to a susceptible check that exhibited infection. These criteria were chosen to reduce false negatives in the bulk as much as possible.

We selected 45 individuals for the 'resistant' bulk and 49 individuals for the 'susceptible' bulk. The QTL-seq analysis was conducted by bulking bam alignment files *in silico* and assaying allele frequency differences genome-wide between the bulks (Figure 2A). Analysis of the bulks indicated one major

QTL located on the proximal end of chromosome 12. There were no other peaks identified, indicating that a single, major locus accounted for smut resistance from Ascasubi. In the following year (2019-2020), we phenotyped a related population of F3:4 progeny with Ascasubi as a parent and Granoleico as the susceptible parent. We selected a set of 48 resistant individuals and the 48 most susceptible individuals. We sequenced those 96 individuals and again analyzed them with QTL-seq analysis. The major locus on chromosome 12 was again indicated as a single, major resistance locus controlling resistance (Figure 2B).

We have identified and validated a major locus controlling smut resistance in peanut. The qualitative inheritance of this trait, its high level of heritability, low environmental variance, and strong phenotyping methodologies, we were able to map and validate this resistance within early generations (F2:3 and F3:4), by phenotyping single plants rather than replicated plots. Phenotyping and genotyping early generation families within a developing population saves time and allows the potential for selection for backcross breeding. This method greatly decreased the time needed for discovery. The resistance is indicated on an approximal 2 Mb region that contains the largest cluster of R genes in the peanut genome (Bertioli *et al.*, 2016). We have generated highly contiguous genome assemblies of the resistant parent, Ascasubi, and of the most widely grown cultivar in Argentina, Granoleico. Granoleico is very susceptible to smut, and the differences between those two genomes within the 2 Mb region will contain the functional variation that confers strong resistance to smut. This resistance and the markers associated with it will be transformational in breeding smut resistant cultivars for Argentina and the world.

Table 1. Assembly stats for the Ascasubi and Granoleico genome assemblies. Contigs were scaffolded into 20 pseudomolecules comprising the 10 A chromosomes and 10 B chromosomes. N/L50 is the length of the contig after half of the assembly has been accounted for when contigs are sorted from largest to smallest. Briefly, for Ascasubi half of the genome is assembled into 16 contigs that are longer than 53 Mb. N/L90 is calculated the same, but for 90% of the assembly. Briefly, for Ascasubi, 90% of the assembly is in 39 contigs which are longer than 19 Mb.

	Ascasubi (R)	Granoleico (S)
Scaffold total	20	20
Contig total	92	97
Contig N/L50	16/53.095 Mb	18/53.926 Mb
Contig N/L90	39/19.502 Mb	39/27.963 Mb
Max contig length	136.375 Mb	159.756 Mb




Figure 1: Smut infection as measured by infected pods/total pods in individual F2:3 progeny plants (A) tested at Manfredi in 2018-2019 field season (B).




Figure 2: Mapping and validation of smut resistance. (A) A major QTL was identified on chromosome 12. Bulk analysis of 45 individuals with low infection and 49 individuals with infection >3% identifies a 2 Mb region conferring resistance. Y axis is the difference in allele frequency between the two bulks. A peak indicates a region where the allele frequencies significantly segregate between the bulks. The X axis is the physical location on the chromosome. (B) Validation of the QTL on chromosome 12 in a validation population of 48 resistant individuals and 48 susceptible individuals.

ACKNOWLEDGEMENTS

Funding for this project was provided by USDA ARS CRIS Projects 3072-21220-008-00D, 6048-21000-029-000D, 6066-21310-005-000D, and 6046-21000-012-000D, the USDA ARS National Plant Disease Recovery System (NPDRS,) and MARS WRIGLEY, Inc. The authors declare no conflict of interest. The authors would like to thank Lisa Myers for technical assistance. Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture. USDA is an equal opportunity provider and employer.

LITERATURE CITED

- Agarwal G., Clevenger J., Pandey M.K., Wang H., Shasidhar Y., Chu Y., Fountain J.C., Choudhary D., Culbreath A.K., Liu X., Huang G., Wang X., Deshmukh R., Holbrook C.C., Bertioli D.J., Ozias-Akins P., Jackson S.A., Varshney R.K., and B. Guo. 2018. High-density genetic map using whole-genome re-sequencing for fine mapping and candidate gene discovery for disease resistance in peanut. *Plant Biotechnol. J.* DOI 10.1111/pbr.1274310.1111/pbi.12930.
- Bertioli D.J., Cannon S.B., Froenicke L., Huang G., Farmer A.D., Cannon E.K.S., Liu X., Gao D., Clevenger J., Dash S., Ren L., Moretzsohn M.C., Shirasawa K., Huang W., Vidigal B., Abernathy B., Chu Y., Niederhuth C.E., Umale, P., Araújo, A.C.G., Kozik, A., Kim, K.D., Burow, M.D., Varshney, R.K., Wang, X., Zhang, X., Barkley, N., Guimarães, Isobe S., Guo B., Stalker H.T., Schmitz R., Scheffler B.E., Leal-Bertioli S.C.M., Xun X., Jackson S.A., Micheltore R., and P. Ozais-Akins. 2016. The genome sequences of *Arachis duranensis* and *Arachis ipaensis*, the diploid ancestors of cultivated peanut. *Nature Genet.* 48:438-446. DOI 10.1038/ng.3517.
- Bertioli D.J., Jenkins J., Clevenger J., Dudchenko O., Gao D., Seijo G., Leal-Bertioli S.C.M., Ren L., Farmer A.D., Pandey M.K., Somoluk S.S., Abernathy B., Agarwal G., Ballén-Taborda C., Cameron C., Campbell J., Chavarro C., Chitikineni A., Chu Y., Dash S., El Baidouri M., Guo B., Huang W., Kim K.D., Korani W., Lanciano S., Lui C.G., Mirouze M., Mortensohn M.C., Pham M., Shin J.H., Shirasawa K., Sinharoy S., Sreedasyam A., Weeks N.T., Zhang X., Zheng Z., Sun Z., Froenicke L., Aiden E.L., Micheltore R., Varshney R.K., Holbrook C.C., Cannon E.K.S., Scheffler B.E., Grimwood J., Ozais-Akins P., Cannon S.B., Jackson S.A., and J. Schmutz. 2019. The genome sequence of segmental allotetraploid peanut *Arachis hypogaea*. *Nature Genet.* 51:877-884. DOI 10.1038/s41588-019-0405-z.
- BICON. 2017. Australian Biosecurity Import Conditions. <https://bicon.agriculture.gov.au/BiconWeb4.0/ViewElement/Alert?elementPk=687363&casePk=682337>.
- Bonessi F., Rago A.M., Marinelli A.D., March G.J., Cazón L.I. García J., and C.M. Oddino. 2011. Efecto de la fertilización sobre la intensidad del carbon del maní. In: *Jornada Nacional del Maní*, General Cabrera, Córdoba, Argentina. pp. 63-64.
- Bressano M., Massa A.N., Arias R.S., de Blas F., Oddino C., Faustini P.C., Soave S., Soave J.H., Pérez M.A., Sobolev V.S., Lamb M.C., Balzarini M., Buteler M.I., and J.G. Seijo. 2019. Introgression of peanut smut resistance from landraces to elite peanut cultivars (*Arachis hypogaea* L.). *PLoS ONE* 14: e0211920. DOI 10.1371/journal.pone.0211920.
- Burow M.D., Simpson C.E., Paterson A.H., and J.L. Starr. 1996. Identification of peanut (*Arachis hypogaea* L.) RAPD markers diagnostic of root-knot nematode (*Meloidogyne arenaria* (Neal) Chitwood) resistance. *Mol. Breed.* 2:369-379. DOI 10.1007/BF00437915.
- Carranza J. M., and J.C. Lindquist. 1962. *Thecaphora frezii* sp., parásita de *Arachis* sp. *Boletín de la Sociedad Argentina de Botánica*, X(1): 11-17.
- Cazón L. I., Conforto C., Paredes J. A., Bisonard E. M., and A.M. Rago. 2014. Sensibilidad y especificidad de la técnica de PCR en la detección de teliosporas de *Thecaphora frezii* en semillas de maní. In: *29° Jornada Nacional de Maní*. General Cabrera, Córdoba, Argentina. pp. 40-42.
- Cazón L.I., Paredes J.A., and A.M. Rago. 2018. The biology of *Thecaphora frezii* smut and its effects on Argentine peanut production. In J.N. Kimatu (ed.) *Advances in plant pathology*, IntechOpen Ltd., London, U.K. pp. 31-46. DOI 10.5772/intechopen.75837.
- Chamberlin K.D., Baldessari J., Bennett R.S., Clevenger J.P., Holbrook C.C., Tallury S.P., Chu Y., Ozias-Akins P., Conde M.B., and M.E. Payton. 2022. Identification of germplasm resistant to peanut smut. *Peanut Sci.* 49: 1-16. DOI 10.3146/0095-3679-491-PS21-10.
- Chenault K.D., Maas A.L., Damicone J.P., Payton M.E., and H.A. Melouk. 2009. Discovery and characterization of a molecular marker for *Sclerotinia minor* (Jagger) resistance in peanut. *Euphytica* 166:357-365.
- Church G.T., Simpson C.E., Burow M.D., Paterson A.H. and J.L. Starr. 2000. Use of RFLP markers for identification of individuals homozygous for resistance to *Meloidogyne arenaria* in peanut. *Nematology* 2:575-580.
- Cignetti M. I., Marraro Acuña F., and P.H. Mazzini. 2010. Influencia de la labranza sobre la intensidad del carbón del maní. Pages 12-14 in: *25 Jornada Nacional del Maní*. General Cabrera, Córdoba, Argentina.
- Clevenger J., Chu Y., Chavarro C., Agarwal G., Bertioli D.J., Leal-Bertioli S.C.M., Pandey M.K., Vaughn J., Abernathy B., Barkley N.A., Hovav R., Burow M., Nayak S.N., Chitikineni A., Isleib T.G., Holbrook C.C., Jackson S.A., Varshney R.K., and P. Ozias-Akins. 2017. Genome-wide

- SNP genotyping resolves signatures of selection and tetrasomic recombination in peanut. *Mol. Plant* 10(2):309–322 DOI 10.1016/j.molp.2016.11.015.
- Clevenger J.P., Korani W., Ozias-Akins P., and S. Jackson. 2018. Haplotype-based genotyping in polyploids. *Front in Plant Sci.* 9:564 DOI 10.3389/fpls.201800564.
- Figueredo M.S., Tonelli M.L., Ibáñez F., Morla F., Cerioni G., Tordable M. C., and A. Fabra. 2017. Induced systemic resistance and symbiotic performance of peanut plants challenged with fungal pathogens and co-inoculated with the biocontrol agent *Bacillus* sp. CHEP5 and *Bradyrhizobium* sp. SEMIA6144. *Microbiol. Res.* 197: 65-73. DOI 10.1016/j.micres.2017.01.002.
- Herselman L., Thwaites R., Kimmins F.M., Courtois B., van der Merwe P.J.A., and S.E. Seal. 2004. Identification and mapping of AFLP markers linked to peanut (*Arachis hypogaea* L.) resistance to the aphid vector of groundnut rosette disease. *Theor. Appl. Genet.* 109:1426-1433.
- Ibáñez M.A., Minudri F.H., Kearney M.I., Rago A.M., Paredes J.A., Mojica C., and E.G. Peiretti. 2018. Análisis multiambiental del comportamiento de genotipos de maní frente a carbón. Proc. "XXXIII Jornada Nacional de Maní", pp. 12. General Cabrera, Córdoba, Argentina. http://ciacabrera.com.ar/jornada_del_maní/33_jornada_del_maní.html.
- Inglis P.W., Pappas Md.C.R., Resende L.V., and D. Grattapaglia. 2018. Fast and inexpensive protocols for consistent extraction of high-quality DNA and RNA from challenging plant and fungal samples for high-throughput SNP genotyping and sequencing applications. *PLoS ONE* 13(10): e0206085. DOI 10.1371/journal.pone.0206085.
- Jadhav Y., Manohar S.S., Sunkad G., Kannalli V.P., Pandey M.K., Variath M.T., Yaduru S., Kona P., Varshney R.K. and J. Pasupuleti. 2019. Genomic regions associated with resistance to peanut bud necrosis disease (PBND) in a recombinant inbred line (RIL) population. *Plant Breed.* 138:748-760. DOI 10.1111/pbr.12743.
- Jasani M.D., Kamdar J.H., Bera S., Sunkad G., and S.K. Bera. 2021. Novel and stable major QTLs conferring resistance to peanut bud necrosis disease and identification of resistant high yielding peanut breeding lines. *Euphytica* 217: 105. DOI 10.1007/s10681-021-02835-7.
- Khan S.A., Chen H., Deng Y., Chen Y., Zhang C., Cai T., Ali N., Mamadou G., Xie D., Guo B., Varshney R.K., and W. Zhuang. 2020. High-density SNP map facilitates fine mapping of QTLs and candidate genes discovery for *Aspergillus flavus* resistance in peanut (*Arachis hypogaea*). *Theor. Appl. Genet.* 133:2239-2257. DOI 10.1007/s00122-020-03594-0.
- Khedikar Y.P., Gowda M.V., Sarvamangala C., Patgar K.V., Upadhyaya H.D., and R.K. Varshney. 2010. A QTL study on late leaf spot and rust revealed one major QTL for molecular breeding for rust resistance in groundnut (*Arachis hypogaea* L.). *Theor. Appl. Genet.* 121:971-84. DOI 10.1007/s00122-010-1366-x.
- Leal-Bertioli S.C.M., Moretzsohn M.C., Roberts P.A., Ballén-Taborda C., Borba T.C., Valdisser P.A., Vianello R.P., Araújo A.C.G., Guimarães P.M., and D.J. Bertioli. 2016. Genetic mapping of resistance to *Meloidogyne arenaria* in *Arachis stenosperma*: a new source of nematode resistance for peanut. *G3: Genes, Genomes, Genet.* 6:377-390. DOI: 10.1534/g3.115.023044.
- Liang Y., Cason J.M., Baring M.R., and E.M. Septiningsih. 2021. Identification of QTLs associated with Sclerotinia blight resistance in peanut (*Arachis hypogaea* L.). *Genet. Resour. Crop Evol.* 68:629-637. DOI 10.1007/s10722-020-01012-4.
- Liao P.Y. and K.H. Lee. 2010. From SNPs to functional polymorphism: the insight into biotechnology applications. *Biochem. Eng. J.* 49:149–158.
- Luo H., Pandey M.K., Zhi Y., Zhang H., Xu S., Guo J., Wu B., Chen H., Ren X., Zhou X., Chen Y., Chen W., Huang L., Liu N., Sudini H.K., Varshey R.K., Lei Y., Liao B., and H. Jiang. 2020a. Discovery of two novel and adjacent QTLs on chromosome B02 controlling resistance against bacterial wilt in peanut variety Zhonghua 6. *Theor. Appl. Genet.* 133:1133-1148. DOI 10.1007/s00122-020-03537-9.
- Luo Z., Cui R., Chavarro C., Tseng Y., Zhou H., Peng Z., Chu Y., Yang Z., Lopez Y., Tillman B., Dufault N., Brenneman T., Isleib T.G., Holbrook C., Ozias-Akins P., and J. Wang. 2020b. Mapping quantitative trait loci (QTLs) and estimating the epistasis controlling stem rot resistance in cultivated peanut (*Arachis hypogaea*). *Theor. Appl. Genet.* 133:1201-1212. DOI 10.1007/s00122-020-03542-y.
- Marinelli A., March G. J., and C. Oddino. 2008. Aspectos biológicos y epidemiológicos del carbón del maní (*Arachis hypogaea* L.) causado por *Thecaphora frezii* Carranza & Lindquist. *AgriScientia*, 25: 1–5. DOI 10.31047/1668.298x.v25.n1.2735.
- Marraro Acuña F. and R.J. Haro. 2011. Carbón del maní (*Thecaphora frezii*): su incidencia en rotaciones de cultivo. In: XXVI Jornada Nacional del Maní, Córdoba, Argentina, pp. 28-30
- Massa A.N., Bressano M., Soave J.H., Buteler M.I., Seijo G., Sobolev V.S., Orner V.A., Oddino C., Soave S.J., Faustinelli P.C., de Blas F.J., Lamb M.C., and R.S. Arias. 2021. Genotyping tools and resources to assess peanut germplasm: smut-resistant landraces as a case study. *Peer J.* DOI 10.7717/peerj.10581.
- Mondal S. and A.M. Badigannavar. 2018. Mapping of a dominant rust resistance gene revealed two R genes around the major Rust_QTL in cultivated peanut (*Arachis hypogaea* L.). *Theor. Appl. Genet.* 131:1671-1681. DOI 10.1007/s00122-018-3106-6.

- Oddino C., Marinelli A., March G., García J., Tarditi L., D'Eramo L., and S. Ferrari. 2010. Relación entre el potencial inóculo de *Thecaphora frezii*. La intensidad de carbón del maní y el rendimiento del cultivo. 25 Jornada de Maní. General Cabrera, Córdoba. CIA-INTA. pp. 24–26.
- Oddino C., Soave J., Soave S., Moresi A., Bianco C., Buteler M., Faustinelli P., and D. Torre. 2013. Avances genéticos en la tolerancia a carbón del maní causado por *Thecaphora frezii*. In: Proc. "XXVIII Jornada Nacional de Maní", pp. 31-32. General Cabrera, Córdoba, Argentina. http://ciacabrera.com.ar/jornada_del_maní/28_jornada_del_maní.html
- Pandey M.K., Agarwal G., Kale S.M., Clevenger J., Nayak S.N., Sriswathi M., Chitikineni A., Chavarro C., Chen X., Upadhyaya H.D., Vishwakarma M.K., Leal-Bertioli S., Liang X., Bertioli D.J., Guo B., Jackson S.A. Ozais-Akins P., and R.K. Varshney. 2017. Development and evaluation of a high-density genotyping 'Axiom_Arachis' array with 58 K SNPs for accelerating genetics and breeding in groundnut. *Sci. Rep.* 7: 40577. DOI 10.1038/srep40577.
- Paredes J.A., Cazón L.I., Oddino C., Monguillot J.H., Rago A.M. and J.P. Edwards Molina. 2021. Efficacy of fungicides against peanut smut in Argentina. *Crop Prot.* 140:105403. DOI 10.1016/j.cropro.2020.105403.
- Rago A.M., Cazón L.I., Paredes J.A., Molina J.P.E., Conforto E.C., Bisonard E.M., and C. Oddino. 2017. Peanut smut: From an emerging disease to an actual threat to Argentine peanut production. *Plant Dis.* 100: 400-408, DOI 10.1094/PDIS-09-16-1248-FE.
- Soave J.H. 2002. Granoleico, nuevo cultivar de maní (*Arachis hypogaea* L.) tipo runner con alta relación oleico-linoleico. In: Proc. "XVII Jornada Nacional de Maní", pp. 40. General Cabrera, Córdoba, Argentina. http://ciacabrera.com.ar/jornada_del_maní/17_jornada_del_maní.html
- Tseng Y., Tillman B.L., Peng Z., and J. Wang. 2016. Identification of major QTLs underlying tomato spotted wild virus resistance in peanut cultivar Florida-EPTM '113'. *BMC Genet.* 17: 1. DOI 10.1186/s12863-016-0435-9.
- Wang H., Pandey M.K., Qiao L., Qin L., Qin H., Culbreath A.K., He G., Varshney R.K., Scully B.T., and B. Guo. 2013. Genetic mapping and qualitative trait loci analysis for disease resistance using F2 and F5 generation-based genetic maps derived from 'Tifrunner' x 'GT-C20' in peanut. *The Plant Gen.* 6:1-10. DOI 10.3835/plantgenome2013.05.0018.
- Wann D.Q., Falco A., Cavigliasso M., and C. Cassano. 2020. Phenotypic variation of peanut smut (*Thecaphora frezii*) incidence and severity in the U.S. peanut mini-core collection. *Peanut Sci.* 47: 46-53. DOI 10.3146/PS20-4.1.
- Workman R., Timp W., Fedak R., Kilburn D., Hao S., and K. Liu. 2018. High Molecular Weight DNA Extraction from Recalcitrant Plant Species for Third Generation Sequencing. Protocol Exchange. DOI 10.1038/protex.2018.059.
- Zhao Z., Tseng Y., Peng Z., Lopez Y., Chen C.Y., Tillman B.L., Dang P., and J. Wang. 2018. Refining a major QTL controlling spotted wild disease resistance in cultivated peanut (*Arachis hypogaea* L.) and evaluating its contribution to the resistance variations in peanut germplasm. *BMC Genet.* 19: 17.