



# REVISTA ARGENTINA DE MICROBIOLOGÍA

[www.elsevier.es/ram](http://www.elsevier.es/ram)



## BRIEF REPORT

### Microscopic and PCR-based detection of microsporidia spores in human stool samples

Cristian Javier Mena<sup>a,b</sup>, Andrés Barnes<sup>b</sup>, Gonzalo Castro<sup>c</sup>, Lorena Guasconi<sup>a</sup>, Verónica Liliana Burstein<sup>a</sup>, Ignacio Beccacece<sup>a</sup>, Patricia Catalina Paulin<sup>d</sup>, Joel Arneodo<sup>e</sup>, Silvana Carnevale<sup>f</sup>, Germán Astudillo<sup>f</sup>, Laura Cervi<sup>a</sup>, Martín Gustavo Theumer<sup>a</sup>, Laura Silvina Chiapello<sup>a,\*</sup>

<sup>a</sup> Departamento de Bioquímica Clínica, Facultad de Ciencias Químicas, Universidad Nacional de Córdoba, Centro de Investigaciones en Bioquímica Clínica e Inmunología, CONICET, Córdoba, Argentina

<sup>b</sup> Sección Parasitología, Laboratorio de Microbiología, Área de Diagnóstico Complementario, Hospital Rawson, Córdoba, Argentina

<sup>c</sup> Área Biología Molecular, Laboratorio Central de la Provincia de Córdoba, Córdoba, Argentina

<sup>d</sup> Laboratorio de Parasitología, Hospital de Pediatría S.A.M.I.C. "Prof. Dr. Juan P. Garrahan", Buenos Aires, Argentina

<sup>e</sup> Instituto de Microbiología y Zoología Agrícola (IMyZA), Instituto Nacional de Tecnología Agropecuaria (INTA), Argentina

<sup>f</sup> Instituto Nacional de Enfermedades Infecciosas, ANLIS "Dr. Carlos G. Malbrán", Buenos Aires, Argentina

Received 30 October 2019; accepted 7 April 2020

#### KEYWORDS

Microsporidiosis;  
*Enterocytozoon bieneusi*;  
Nested PCR;  
Trichrome stain

**Abstract** Microsporidia are obligate intracellular fungi with a remarkable ability to infect a wide range of invertebrate and vertebrate hosts. Namely, *Enterocytozoon bieneusi* is the most frequently microsporidia reported worldwide, and mainly associated with chronic diarrhea and wasting syndrome in AIDS patients. Microscopy and PCR-based detection techniques are effective for diagnosis and identification of species and genotypes; however, these methods should be standardized in each laboratory. In this study, we performed microscopy and nested PCR techniques with PCR product sequencing to detect *E. bieneusi* in human stool samples. These techniques, if applied together, might prove useful for diagnosis and future epidemiological studies of intestinal microsporidiosis in Argentina.

© 2020 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

#### PALABRAS CLAVE

Microsporidiosis;  
*Enterocytozoon bieneusi*;

Detección microscópica y molecular de esporas de microsporidios en muestras de heces humanas

**Resumen** Los microsporidios son hongos intracelulares obligados con una notable capacidad para infectar una amplia gama de hospedadores invertebrados y vertebrados. *Enterocytozoon bieneusi* es el microsporidio más frecuentemente reportado en todo el mundo, principalmente

\* Corresponding author.

E-mail address: [laura.chiapello@unc.edu.ar](mailto:laura.chiapello@unc.edu.ar) (L.S. Chiapello).

<https://doi.org/10.1016/j.ram.2020.04.005>

0325-7541/© 2020 Asociación Argentina de Microbiología. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PCR anidada;  
Coloración tricrómica

asociado con diarrea crónica y síndrome debilitante en pacientes con sida. Las técnicas de detección basadas en microscopía y PCR son útiles para el diagnóstico y la identificación de especies y genotipos, pero estos métodos deben estar estandarizados en cada laboratorio. En este estudio evaluamos técnicas de microscopía y PCR anidada, con secuenciación de los productos, para detectar *E. bieneusi* en muestras de heces humanas. Estas técnicas, usadas conjuntamente, podrían ser útiles para su aplicación en el diagnóstico de microsporidiosis intestinal y para realizar estudios epidemiológicos de esta afección en Argentina.

© 2020 Asociación Argentina de Microbiología. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Microsporidia are obligate intracellular parasites with a remarkable ability to exploit a wide range of invertebrate and vertebrate animals, causing from cryptic, benign infections to massive infections with extensive damage leading even to host death<sup>5</sup>. These organisms were previously considered "primitive" protozoa; however, molecular phylogenetic studies as well as biological evidence have demonstrated that microsporidia are related to the kingdom Fungi, either as a basal branch or sister group<sup>5</sup>. Although there are 8 genera and 14 species able to infect humans, two species are the most commonly identified, *Enterocytozoon bieneusi* and *Encephalitozoon intestinalis*. Human infections are thought to be mainly zoonotic or waterborne and microsporidia spores are released into the environment with feces, urine, and respiratory secretions<sup>5</sup>. Immuno-competent individuals have asymptomatic or self-limited infections, occasionally reported as "traveler's diarrhea"; however, AIDS patients or other immunosuppressed individuals suffer intestinal microsporidiosis with chronic diarrhea, malabsorption syndrome or severe disseminated disease<sup>5</sup>. The prevalence of this emerging opportunistic infection in human ranges from 1 to over 50%, depending on the geographic region, the methods applied for diagnosis and demographic characteristics of the studied population<sup>11</sup>. Albendazole is effective against many species including *Encephalitozoon* spp., but it shows low efficacy in controlling *E. bieneusi* disease<sup>5</sup>.

Transmission electron microscopy (TEM) is regarded as the golden standard for establishing the diagnosis but is expensive, time-consuming and has low sensitivity. Light or immunofluorescence microscopy and molecular amplification techniques, primarily using PCR, are now routinely used for diagnosis. As a matter of fact, PCR assays are significantly more sensitive than light microscopy and provide species differentiation, thus having an impact on treatment decisions<sup>3</sup>.


In Argentina, 120 000 people live with *Human immunodeficiency virus* (HIV)/acquired immune deficiency syndrome (AIDS) and 6500 new cases are reported annually. In turn, 27.5% of these people are diagnosed late; therefore, they are often at risk of acquiring an opportunistic infection<sup>7</sup>. Nevertheless, there are few epidemiological data on microsporidiosis<sup>13</sup>, and a reliable diagnosis is particularly necessary for AIDS patients with chronic diarrhea<sup>12</sup>. In this study, we standardized microscopy techniques as well as a

nested PCR to detect microsporidia in stool samples<sup>2,8</sup> so as to be further applied in diagnosis and epidemiological studies in public health in Argentina.

Stool samples (Sample A and B, non-HIV) from Rawson Hospital (Córdoba, Argentina) stock collection samples were used as negative controls and the *E. bieneusi* positive sample was provided by the Parasitology Laboratory of Hospital Garrahan, Buenos Aires, Argentina. The research protocol was approved by the Ethical Committee for Research of Rawson Hospital, Córdoba, Argentina (number 05072018).

Chromotrope 2R technique was performed by following the method described by Weber et al.<sup>14</sup>, with slight modifications. Briefly, stool samples were fixed with ethanol 70% (1/3 ratio) and then concentrated by the modified ethyl-acetate stool-concentration method<sup>4</sup>. Smears were fixed with methanol for 5 min and incubated overnight with chromotrope 2R stain, prepared as described by Weber et al.<sup>14</sup>. After staining, slides were rinsed and de-colored by dipping in acid alcohol for 10 s (moving the slide up and down) and stopped with 95% alcohol. Smears were successively dehydrated in 95% alcohol (5 min), 100% alcohol (10 min) and xylene (10 min). In parallel smears, calcofluor white staining was also performed as previously described<sup>6</sup>.

To confirm microsporidia detection by light microscopy, we further performed a nested technique as previously described<sup>1</sup> and modified by Santin-Duran et al.<sup>2,9,10</sup>. DNA extraction was performed by using the DNAeasy Blood & Tissue Kit (Qiagen, USA), according to the manufacturer's instructions. PCR was performed using a set of nested primers specific for *E. bieneusi* that amplified the ITS and portions of the flanking large and small subunits of the rDNA (~400 bp). The outer primers were EBITS3 (5'-GGTCATAGGGATGAAGAG-3') and EBITS4 (5'-TTCGAGTTCTTCGCGCTC-3'), and the inner primers were EBITS1 (5'-GCTCTGAATATCTATGGCT-3') and EBITS2 (5'-ATCGCCGACGGATCCAAGTG-3')<sup>2</sup>. The reaction mixture (20 µl) contained 2 µl of buffer 10× (16 mM MgCl<sub>2</sub>, 500 mM KCl, 100 mM Tris-HCl; Dongsheng Biotech, China); 0.16 µl of MgCl<sub>2</sub> 50 mM; 0.4 µl of 10 mM dNTPs; 20 pmol of each primer; 0.2 µl of 5 U/µl of Taq polymerase (Dongsheng Biotech); and 5 µl of purified DNA. After denaturation at 94 °C for 3 min, the PCR samples were subjected to 35 cycles of amplification (denaturation at 94 °C for 30 s, annealing at 57 °C for 30 s, and elongation at 72 °C for 40 s), followed by a final extension at 72 °C for 10 min. Nested PCR conditions were




**Figure 1** Weber's modified trichome (A and B) and calcofluor (C) staining showing *Enterocytozoon bieneusi* spores in feces. (A) Modified trichrome staining observed by light microscopy showing ovoid shaped-spores with a pinkish-red stained wall (arrows). Some spores show a pinkish belt-like stripe. (B) Calcofluor staining observed by fluorescence microscopy showing positive *E. bieneusi* spores (arrows) and a yeast (asterisk). Magnification 1000 $\times$ . Bars = 5  $\mu$ m.

identical to those of the first run except that only 30 cycles were carried out with an annealing temperature of 55 °C. Pure and diluted (1/10, 1/100 and 1/100) PCR products were used in the nested PCR. A negative control without DNA was included in all PCR sets. Furthermore, to rule out PCR inhibition, DNA amplification from *Candida albicans* was also performed in all samples (Supplementary Fig.). PCR products were subjected to electrophoresis in 2% agarose gel and visualized by staining the gel with ethidium bromide.

To determine *E. bieneusi* identity, the amplification product from the nested PCR was then subjected to a direct nucleotide sequencing reaction in both directions by using the inner-nested PCR primers (EBITS1 and EBITS2) in the Applied Biosystems® 3500xL Genetic Analyzer at Laboratorio Central de la Provincia de Córdoba, Argentina. Sequences obtained were aligned and inspected by using the MEGA software (version 6), and then compared with reference sequences from the GenBank database by the BLAST analysis<sup>10</sup>.

Microsporidia are still difficult to diagnose even though significant progress has been made over the last few years. Nowadays microscopic or PCR-based detection techniques,

which must be standardized in each laboratory<sup>4</sup>, are useful for diagnosis and genotyping. In this study, we standardized the most commonly used staining technique for the examination of stool specimens: chromotrope 2R-based (Weber's modified trichrome) stain<sup>14</sup> and chemiluminescent optical brightening agent calcofluor white (that detects chitin in the spore cell wall)<sup>4</sup>. Figure 1A shows pinkish red-stained microsporidia spores, with some of them showing a distinct pinkish belt-like stripe (1000 $\times$ ). The spores show a characteristic staining pattern and can be readily distinguished from bacteria and fecal debris<sup>4,14</sup>. In addition, as described by Weber et al.<sup>14</sup>, calcofluor stained not only microsporidia spores but also yeasts and some feces debris (Figure 1B). In this regard, previous studies comparing the chromotrope staining technique with methods using chemiluminescent optical brighteners have shown that these tests are robust for routine use<sup>4</sup>. However, none of these microscopic methods are sufficient to determine microsporidia species; therefore, we also performed a nested PCR that is currently used to identify *E. bieneusi* genotypes and other microsporidia from stool samples. Figure 2 shows that nested PCR produced fragments of approximately 390 bp



**Figure 2** Agarose gel electrophoresis (2%) showing amplification products from the nested PCR from ethanol-fixed stool samples. (A) PCR from *Enterocytozoon bieneusi* positive stool sample. L: molecular weight ladder. Lanes 1–3: products of nested PCR (with inner primers ITS1 and ITS2) using diluted products from PCR as DNA template (Lane 1: 1/10, Lane 2: 1/100, Lane 3: 1/1000). Lane 4: PCR product using undiluted DNA template from PCR. (B) Nested PCR from different stool samples. L: molecular weight ladder, Lane 1: negative control ( $H_2O$ ), Lane 2: negative stool sample A (PCR product diluted 1/100), Lane 3: negative stool sample B (1/100), Lane 4: positive stool sample (1/100), Lane 5: negative stool sample A (diluted 1/10), negative stool sample B (diluted 1/10).

(Fig. 2A and B). Furthermore, the best performance was obtained when products from the first amplification reaction (435 bp, not shown) were diluted (1/100 and 1/1000) to be used in the nested PCR (Fig. 2A).

On the other hand, this PCR technique amplifies the internal transcribed spacer (ITS) region of the rRNA, which has been useful in many studies for the identification of *E. bieneusi* genotypes<sup>2,3,8–10</sup>. Accordingly, the sequence analysis of PCR-amplified products showed 99.68% (319 pb) homology with genotype B of *E. bieneusi* (GenBank accession number AF101198.1)<sup>10</sup>.

In conclusion, in this work we successfully detected *E. bieneusi* microsporidia from a human stool sample by using the Chromotrope 2R microscopic technique and a previously reported nested PCR, which can also be applied for *E. bieneusi* genotyping<sup>2,8–10</sup>. Further studies are currently underway to define whether these techniques could be useful for human microsporidiosis surveillance in Argentina.

## Funding

This work was supported by the Secretary of Science and Technology of the National University of Córdoba (Sec-Subsecretaría de Ciencia y Tecnología de la Universidad Nacional de Córdoba, Argentina, SECyT-UNC, PROYECTO CONSOLIDAR N°33620180100408) and the National Agency for Scientific and Technological Promotion – Fund for Scientific and Technological Research (Agencia Nacional de Promoción Científica y Tecnológica, Fondo para la Investigación Científica y Tecnológica, Argentina, FONCyT, PICT 2015-1425). CJM is a fellow from SECyT-UNC (Grants for Socio-productive Technological Innovation - Becas de Innovación Tecnológica Socio-Productiva, BITS 2018) and LSC is a member of the scientific career of the National Council for Scientific and Technological Research (Consejo Nacional de Investigaciones Científicas y Tecnológicas, CONICET), Argentina.

## Conflict of interest

The authors declare that they have no conflicts of interest.

## Acknowledgment

We wish to thank Dr. Mónica Santin-Duran (United States Department of Agriculture, Maryland, USA) for providing PCR protocols and Dr. Sebastián Dambolena (IMVIB-CONICET, Córdoba, Argentina) for technical support.

## Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.ram.2020.04.005

## References

1. Buckholt M, Lee J, Tzipori S. Prevalence of *Enterocytozoon bieneusi* in swine: an 18-month survey at a slaughterhouse in Massachusetts. Appl Environ Microbiol. 2002;68:2595–9.
2. Del Coco V, Córdoba M, Bilbao G, de Almeida Castro P, Basualdo J, Santin M. First report of *Enterocytozoon bieneusi* from dairy cattle in Argentina. Vet Parasitol. 2014;199:112–5.
3. Ghosh K, Weiss L. Molecular diagnostic tests for Microsporidia. *Interdiscip Perspect Infect Dis*. 2009;926521:1–13.
4. Ghosh K, Schwartz D, Weiss LM. Laboratory diagnosis of Microsporidia. In: Weiss LM, Becnel JJ, editors. *Microsporidia: pathogens of opportunity*. 1st ed. Iowa: Wiley; 2014. p. 421–56.
5. Han B, Weiss L. Microsporidia: obligate intracellular pathogens within the fungal kingdom. *Microbiol Spectr*. 2017;5:1–17.
6. Ramanan P, Pritt BS. Extraintestinal microsporidiosis. *J Clin Microbiol*. 2014;52:3839–44.
7. Riera F, Caeiro J, Denning D. Burden of serious fungal infections in Argentina. *J Fungi (Basel)*. 2018;4:1–11.
8. Santin M, Calero-Bernal R, Carmena D, Mateo M, Balseiro A, Barra M, Lima Barbero JF, Habela MA. Molecular characterization

Microscopic and PCR-based detection of microsporidia spores in human stool samples

5

- of *Enterocytozoon bieneusi* in wild carnivores in Spain. *J Eukaryot Microbiol.* 2018;65:468–74.
9. Santin M, Fayer R. Microsporidiosis: *Enterocytozoon bieneusi* in domesticated and wild animals. *Res Vet Sci.* 2011;90:363–71.
10. Santin M, Trout J, Fayer R. *Enterocytozoon bieneusi* genotypes in dairy cattle in the eastern United States. *Parasitol Res.* 2005;97:535–8.
11. Stentiford G, Becnel J, Weiss L, Keeling P, Didier E, Williams B, Bjornson S, Kent M, Freeman M, Brown M, Troemel E, Roesel K, Sokolova Y, Snowden K, Solter L. Microsporidia – emergent pathogens in the global food chain. *Trends Parasitol.* 2016;32:336–48.
12. Velasquez J, Carnevale S, Guarnera E, Labbe J, Chertcoff A, Cabrera M, Rodriguez M. Detection of the microsporidian parasite *Enterocytozoon bieneusi* in specimens from patients with AIDS by PCR. *J Clin Microbiol.* 1996;34:3230–2.
13. Velasquez J, di Risio C, Etchart C, Chertcoff A, Astudillo O, Carnevale S. Multimethodological approach to gastrointestinal microsporidiosis in HIV-infected patients. *Acta Parasitol.* 2019;64:658–69.
14. Weber R, Bryan R, Owen R, Wilcox C, Gorelkin L, Visvesvara G. Improved light-microscopical detection of microsporidia spores in stool and duodenal aspirates. *N Engl J Med.* 1992;326:161–6.