

Crops and Soils Research Paper

Cite this article: Blajman JE, Vinderola G, Páez RB, Signorini ML (2020). The role of homofermentative and heterofermentative lactic acid bacteria for alfalfa silage: a meta-analysis. *The Journal of Agricultural Science* 1–12. <https://doi.org/10.1017/S0021859620000386>

Received: 28 November 2019

Revised: 1 April 2020

Accepted: 21 April 2020

Key words:

Ensiling; *Enterococcus faecium*; *Lactobacillus buchneri*; *Lactobacillus plantarum*; Lucerne; *Pediococcus acidilactici*

Author for correspondence:

J. E. Blajman,

E-mail: jblajman@yahoo.com.ar

The role of homofermentative and heterofermentative lactic acid bacteria for alfalfa silage: a meta-analysis

J. E. Blajman¹ , G. Vinderola², R. B. Páez³ and M. L. Signorini¹

¹Consejo Nacional de Investigaciones Científicas y Técnicas, Instituto Nacional de Tecnología Agropecuaria EEA Rafaela, Ruta 34 Km. 227, 2300 Rafaela, Santa Fe, Argentina; ²Instituto de Lactología Industrial (CONICET-UNL), Facultad de Ingeniería Química, Universidad Nacional del Litoral, Santiago del Estero 2829, 3000 Santa Fe, Argentina and ³Instituto Nacional de Tecnología Agropecuaria EEA Rafaela, Ruta 34 Km. 227, 2300 Rafaela, Santa Fe, Argentina

Abstract

Lactic acid bacteria (LAB) are usually employed as alfalfa silage inoculants to obtain high-quality feed for animal husbandry. However, the effects of these inoculants are still unclear and need to be studied extensively. Therefore, the objective of this meta-analysis was to quantitatively summarize published research studies that assess the effects of homofermentative (HoLAB) and heterofermentative lactic acid bacteria (HeLAB) on fermentation parameters, nutritive value, microbiological composition and aerobic stability of alfalfa silage. PubMed, ScienceDirect and Scopus have been screened for articles published from 1980 to 2018. The criteria for inclusion were: randomized and controlled trials using alfalfa silage and published in peer-reviewed journals. It was found that inoculation with LAB decreased silage pH, neutral detergent fibre, acid detergent fibre and ammoniacal nitrogen, while it increased dry matter and crude protein compared to control in the pooled raw mean difference random-effect model. Additionally, LAB inoculation decreased acetate, propionate, ethanol and butyrate concentrations, whereas it increased lactate. In addition, inoculants reduced the counts of yeasts and moulds. Lastly, LAB inoculation improved aerobic stability. To the best of our knowledge, this is the first meta-analysis that aims at comparing the application of HoLAB and HeLAB for alfalfa silage. In the pool estimate, positive effects attributable to the application of microbial silage inoculants were found in most of the evaluated parameters; supporting the importance of applying both types of inoculants to improve forage preservation for the livestock industry.

Introduction

Alfalfa (*Medicago sativa* L.) has one of the highest feeding values of leguminous forages (Zhang *et al.*, 2017), and is an excellent feed ingredient preserved as hay, haylage or silage for lactating dairy cows (Schmidt *et al.*, 2009). In comparison with grasses, it has a higher content of crude protein (CP), greater concentration of organic acids and minerals; however, it is difficult to ensile due to its high buffering capacity for acidic conditions, in combination with the low water-soluble carbohydrates (WSC) concentrations (McDonald *et al.*, 1991; Ogunade *et al.*, 2016).

The utilization of microbial additives for the purpose of achieving a proper fermentation and improving digestibility has generated great interest, which is evidenced by the fact that inoculants have been developed as silage additives for over 40 years (McDonald *et al.*, 1991; Dunière *et al.*, 2013). These products included strains of homofermentative lactic acid bacteria (HoLAB) or heterofermentative lactic acid bacteria (HeLAB) (Muck, 2010). Administration of HoLAB often boosts silage fermentation by transforming available sugars into lactic acid, thus accelerating the rate of decrease in pH (Contreras-Govea *et al.*, 2011). It should be pointed out that HoLAB inoculants are generally preferred for legume silages as they minimize dry matter (DM) losses through a higher lactic acid production. On the other hand, HeLAB species produce lactic acid and carbon dioxide, as well as traces of ethanol or acetic acid as by-products (Borreani *et al.*, 2018). This leads to DM losses associated with gas production and reduces the feeding value of silage (Ni *et al.*, 2015; Borreani *et al.*, 2018). However, HeLAB inoculants are valuable in enhancing aerobic stability since moderate acetic acid production has the potential to inhibit yeasts and moulds, responsible for initiating spoilage upon exposure to air (Muck *et al.*, 2018; Ferrero *et al.*, 2019). For this reason, modest DM losses from HeLAB treatment should be compensated with improvements in aerobic stability and reduced losses at feed out (Borreani *et al.*, 2018). As both LAB types take different approaches to direct fermentation in the silo, their combination could have potential advantages and complementary effects (Zhang *et al.*, 2009).

Even though many authors have reported several benefits of LAB inoculation on fermentation patterns, there are still unanswered questions and challenges about the extent of variability in the effects of inoculants on the preservation of silage and the impact of the interactions between inoculants and other covariates (for instance, LAB species, LAB application rate, concomitant use of enzymes, study duration and silo scale). Some studies have shown that inoculants enhance the attributes of silages (Zielińska *et al.*, 2015; Liu *et al.*, 2016; Li *et al.*, 2018), but in others experiments, LAB did not consistently decrease ammonia nitrogen production (NH₃-N) (Kozelov *et al.*, 2008), did not preserve DM and CP content (Chilson *et al.*, 2016), failed to improve feed efficiency (Rabelo *et al.*, 2018) and inhibit undesirable microorganisms (Twarużek *et al.*, 2016). Disagreements in responses to LAB addition are multifactorial, and can possibly be attributed to essential factors such as forage maturity, harvesting conditions, moisture content, silage density, mode and application rate of LAB, epiphytic LAB population, sugar availability, plant DM concentration, ensiling duration, efficacy of the inoculant strains and interactions between microbial species in the inoculant and chemical components within the forage (Santos and Kung, 2016; Ozduven and Celebicam, 2017). Additional factors that may explain the reported variability include using experiments with insufficient statistical power and inappropriate experimental designs (Arriola *et al.*, 2017).

A meta-analysis is a highly valuable statistical tool whose objective is to summarize, integrate and contrast the results of a large number of primary studies that investigate the same topic (Shelby and Vaske, 2008). As a result, the meta-analysis generates a more accurate estimate of the effect size of a particular event with greater statistical power than if only one single study was considered (Borenstein *et al.*, 2009). With this perspective in mind, the objective of this work has been to quantitatively summarize published research studies so as to evaluate the magnitude of effects of HoLAB and HeLAB on fermentation parameters, nutritive value, microbiological composition, as well as the outcomes on aerobic stability of alfalfa silage.

Materials and methods

Search strategy

PubMed, ScienceDirect and Scopus databases were screened for articles restricted by language (English, Spanish and Portuguese). The studies included in this meta-analysis were selected only if they were randomized and controlled trials using alfalfa silage, and results were published in peer-reviewed journals between 1980 and 19 April 2018. To evaluate the effects of applying LAB inoculants on fermentation parameters, nutritive value, microbiological composition and aerobic stability of alfalfa silage, peer-reviewed manuscripts were retrieved using the terms 'silage', 'alfalfa', 'lucerne' and 'inoculant'. Studies must have examined uninoculated and inoculated treatment groups, held treatments comprising only LAB and reported response variables with the measures of variance (standard deviation, standard error or variation coefficient). Reviews, duplicate reports, experiments that used different forage species and a number of studies that evaluated other additives were excluded. The term 'study' refers to a scientific article, which can involve one or more experiments. Preliminary screening of titles and abstracts was carried out for eligibility to this study according to the inclusion and exclusion criteria.

Outcomes and definitions

Supplementation with LAB was analysed as a tool which may improve fermentation parameters, nutritive value, microbiological composition and aerobic stability of alfalfa silage. In all studies, the same method was used to measure aerobic stability, which was defined as the number of hours that silage remained stable before increasing more than 2°C above the ambient temperature (Kung and Ranjit, 2001). When the study included more than one inoculant, or when different doses of the same inoculant were used, each inoculated group was compared with the uninoculated group separately.

Data extraction

Information on study design, the number of replicates, means and variances was extracted from each research report. Data for pH, DM concentration (g/kg), neutral detergent fibre (NDF), acid detergent fibre (ADF), NH₃-N (g/kg total N), CP, WSC, ash, ethanol, lactate, acetate, propionate and butyrate, *in vitro* DM digestibility-48 h (IVDMD-48 h), counts of LAB, yeasts, moulds (log₁₀ cfu/g) and aerobic stability (h) were used to estimate outcomes. Certain response variables (DM recovery, lignin, acid detergent-insoluble nitrogen, counts of clostridia and mycotoxins) were retained in the analysis as there were relatively few comparisons that met our selection criteria. For each study, the methodology employed to achieve the results was assessed in detail. However, no scores were used to exclude studies (Lean *et al.*, 2009).

Statistical analysis

The statistical analysis was conducted in Comprehensive Meta Analysis version 2.2 (2011). Due to continuous variables being analysed, results were evaluated by examining the raw mean differences between the inoculant treatment and controls with 95% confidence intervals using a random-effects model. In this model, the true effect may vary from experiment to experiment; we have included between-experiment variability (true heterogeneity) as well as sampling error (Borenstein *et al.*, 2009). To account for variation in precision across studies, the inverse of the squared standard error of each treatment mean was used as a factor in the weight statement of the model.

Heterogeneity, meta-regression and publication bias

A meta-regression analysis was performed to examine heterogeneity sources in the treatment effects. Meta-regression allowed assessing the relationship between year of publication, application rate of LAB inoculant (which ranged from 4 to 7 log₁₀ cfu/g) and duration of the studies as covariates, and silage attributes as outcome variables.

A priori sub-group analyses were planned depending on factors that could potentially influence the magnitude of the treatment: (1) for the purpose of grouping the newest articles, we used the last 10 years (before 2009 *v.* after 2009) as a pre-specified cut-off; (2) type of inoculum (mono-strain *v.* multi-strain); (3) among mono-strain inoculum, type of LAB (HoLAB *v.* HeLAB); (4) LAB species used (with *Lactobacillus buchneri*, with *L. plantarum*, with *Pediococcus acidilactici* and with *Enterococcus faecium*); (5) enzymatic additives addition (with fibrolytic enzymes *v.* without fibrolytic enzymes); (6) study duration (from 30 to 60 days *v.* more than 60 days); and (7) silo type (laboratory or farm scale).

Heterogeneity among studies was assessed using the DerSimonian and the Laird test (Q -statistic). The degree of heterogeneity was quantified with the Inconsistency index (I^2 -statistic; Higgins and Thompson, 2002). An adjusted rank correlation test using the Egger method (Egger *et al.*, 1997) and the Begg test (Begg and Mazumdar, 1994) was used to assess publication bias. It was considered that there was bias if both statistical methods were significant ($P < 0.01$). When there was any evidence of publication bias, the 'trim' and 'fill' method (Duval and Tweedie, 2000) was applied to estimate the quantity and magnitude of missing studies and resultant unbiased effect size. Significance was declared at $P \leq 0.05$ and tendencies at $0.05 > P > 0.1$.

Results

Excluded studies

The literature search yielded 2173 scientific articles on alfalfa silage inoculants. Of the studies identified at the beginning of the meta-analysis, 1976 were excluded on the basis of publication type: articles involving other forage species or mixed crops ($n = 429$), other additives ($n = 88$), or both of them ($n = 265$), reviews ($n = 89$), duplicate reports ($n = 593$), wrong topics ($n = 505$) and studies using inoculants for other purposes ($n = 7$) were rejected. In addition, experiments which were eligible for quantitative review were exempted due to lack of statistical information for conducting a meta-analysis ($n = 25$), studies conducted to assess the impact of certain pathogens like *Escherichia coli* ($n = 2$), manuscripts involving hay or haylage ($n = 12$), papers using simulation models ($n = 3$), no full-text articles ($n = 12$), no English, Spanish or Portuguese language full articles ($n = 3$), studies that analysed the efficacy of symbiotics ($n = 1$), yeasts or propionic acid bacteria ($n = 3$), books or book chapters ($n = 46$) and summaries ($n = 42$) (Fig. 1).

Overview of included studies

At the end of the literature review, 48 studies (131 experiments) were included in this meta-analysis to estimate the role of HoLAB and HeLAB for alfalfa silage. Most of the research papers reviewed did not assess the LAB inoculants' effect over all the parameters under study. Consequently, the number of studies included in the meta-analysis differed in each variable considered. Of the screened experiments, 58 were published before 2009 and the remaining 73 after 2009. Fifty-nine experiments included mono-strain inoculum and 71 included multi-strain LAB. Among mono-strain inoculum, 50 experiments were carried out using HoLAB, whereas nine utilized HeLAB. A total of 18 studies used *L. buchneri* (nine alone and nine in combination with other LAB), 82 used *L. plantarum* (31 alone and 51 in combination with other LAB), 27 used *P. acidilactici* (two alone and 25 in combination with other LAB) and 26 used *E. faecium* (one alone and 24 in combination with other LAB). Inoculants were incorporated with enzymes (20) or without enzymes (111). Studies were conducted for ≤ 60 days (76), or for > 60 days (52). In most of the experiments (118), the inoculant was employed in laboratory-scale silos, while 13 studies were executed in farm-scale silos.

Alfalfa silage conservation

The effects of LAB inoculation on silage quality across studies are depicted in Table 1. In the pooled estimate, inoculation with LAB

decreased silage pH, NDF, ADF and $\text{NH}_3\text{-N}$, whereas DM and CP were increased compared to controls. In contrast, there were no statistical differences in WSC, IVDMD-48 h and ash. Additionally, LAB inoculation reduced acetate, propionate, ethanol and butyrate concentrations, whereas increased lactate. Moreover, LAB inoculation reduced the counts of yeasts and moulds, but did not alter LAB counts. Finally, LAB inoculation improved aerobic stability (Table 1).

Significant heterogeneity (I^2 statistic $> 50\%$) was observed across all silage quality response variables, except for aerobic stability ($I^2 = 41.3\%$). Hence, sub-groups were evaluated in order to identify the sources of variability. In accordance with the sub-group analysis, and only considering significant variables in the pool estimate, inoculation decreased pH in all conditions ($P < 0.001$), except when enzymes were applied ($P = 0.957$) (Tables 2 and 3).

Dry matter increased significantly with inoculation when studies were shorter than 60 days ($P < 0.001$), in those experiments in which inoculants were applied to mini silos ($P < 0.001$) and by HoLAB when mono-strain inoculums were used ($P < 0.001$). Silage inoculation with LAB significantly decreased NDF ($P < 0.001$). Nevertheless, inoculants had no effects when HeLAB were used ($P = 0.320$). Moreover, inoculation decreased ADF ($P < 0.001$). This effect was observed in studies that used mono-strain inoculants ($P < 0.001$) and HoLAB ($P < 0.001$), and in mini silos ($P < 0.001$) (Tables 2 and 3).

Inoculants improved silage protein preservation ($P < 0.001$) in the experiments carried out after 2009 ($P < 0.001$), in the absence of enzymes ($P < 0.001$), by mono-strain LAB ($P < 0.001$) and by HoLAB when single-strain inoculums were used ($P < 0.001$). LAB reduced ($P < 0.001$) $\text{NH}_3\text{-N}$ concentrations in the pool estimate and in the sub-group analysis, with the exception of studies that reported concomitant use of enzymes ($P = 0.286$) (Tables 2 and 3).

With respect to organic acids, ethanol concentrations considerably decreased in all conditions ($P < 0.001$), except when HeLAB ($P = 0.202$) or enzymes ($P = 0.294$) were used. In contrast, lactate significantly increased ($P < 0.001$) when LAB were inoculated, and effects were independent of the sub-group considered. Inoculation with LAB decreased acetate concentrations ($P < 0.001$), but HeLAB significantly increased this organic acid ($P < 0.001$). Propionate decreased in studies conducted before 2009 ($P < 0.001$) and in the absence of enzymes ($P < 0.001$), when LAB were inoculated in laboratory-scale silos ($P < 0.001$), with a multi-strain inoculum ($P < 0.001$), and in experiments shorter than 60 days ($P < 0.001$). The positive effect on butyrate was observed when the experiments were performed by mono-strain LAB ($P < 0.001$) and HoLAB ($P < 0.001$), in the absence of enzymes ($P < 0.001$) and when studies were executed for more than 60 days ($P < 0.001$) (Tables 2 and 3).

Regarding microbiological composition, the number of yeast counts was reduced ($P < 0.001$) when inoculants were used. Conversely, they increased in studies conducted before 2009 ($P < 0.001$) and in those carried out for less than 60 days ($P < 0.001$). No effects were observed on yeast counts with multi-strain inoculum ($P = 0.735$) and with the use of enzymes ($P = 0.094$). The inoculation of LAB showed a positive impact on moulds counts ($P = 0.002$), but the reduction was not significant with the use of enzymes ($P = 0.105$) (Tables 2 and 3).

There were no significant differences regarding the use of HeLAB, simultaneous use of enzymes and long storage periods, which can be attributed to the small number of comparisons found. The low number of studies that incorporate these covariates limited our ability to detect significant effects.




Fig. 1. Flow diagram of studies selected for meta-analysis in this work.

Finally, regarding aerobic stability, this parameter was higher in inoculated alfalfa silage ($P < 0.001$) (Table 1). Nevertheless, it was not possible to conduct a sub-group analysis because all the studies were conducted after 2009, with multi-strain inoculum, without the administration of enzymes, for less than 60 days and in mini silos.

Inoculation with HoLAB (which included *L. plantarum*, *P. acidilactici* and *E. faecium* among others, with the prevalence of the first one) or HeLAB (with *L. buchneri* as the predominant species) showed opposite behaviour in the following response variables: DM ($P = 0.015$), WSC ($P < 0.001$) and acetate concentrations ($P < 0.001$). While HoLAB increased DM and decreased the concentrations of WSC and acetate, HeLAB reduced DM and increased WSC and acetate concentrations (Table 2).

Considering the LAB species included, the absence of *L. buchneri* decreased pH, NDF, ADF, acetate and propionate concentrations ($P < 0.05$), while increased DM and lactate ($P < 0.001$).

Inoculation with *L. buchneri* was effective in decreasing the number of yeasts ($P < 0.001$). CP increased ($P < 0.05$), while $\text{NH}_3\text{-N}$, ethanol, butyrate and moulds counts ($P < 0.05$) were reduced both in the presence and in the absence of *L. buchneri* in the inoculated group (Table 2).

Treating alfalfa silage with *L. plantarum* increased DM concentration ($P < 0.001$). This microorganism was also able to induce a reduction in ethanol ($P < 0.001$), propionate ($P < 0.001$) and acetate concentrations ($P < 0.001$). The absence of *L. plantarum* produced an increase in CP ($P < 0.001$). In addition, a reduction of yeast counts was observed in the inoculated group in the absence of this species ($P < 0.001$). Other measured parameters such as pH, NDF, ADF, $\text{NH}_3\text{-N}$, butyrate and moulds counts decreased ($P < 0.05$), while lactate increased both in the presence and in the absence of *L. plantarum* in the inoculated group ($P < 0.001$) (Table 2).

A significant increase in DM and a reduction in $\text{NH}_3\text{-N}$ and propionate ($P < 0.05$) were observed in the inoculated group in

Table 1. Effects of LAB inoculants on fermentation parameters, nutritive value, microbiological composition and aerobic stability of alfalfa silage (g/kg DM, unless otherwise stated)

Response variable	Control mean (s.e.)	Inoculated mean (s.e.)	No. of trials	RMD	Lower limit	Upper limit	P value	I ² (%)
pH	5 ± 0.05	4.7 ± 0.05	106	-0.3	-0.397	-0.290	<0.001	99.509
DM	335 ± 5.75	340.9 ± 6.19	88	5.9	3.92	7.78	<0.001	96.039
NDF	387 ± 7.27	374.8 ± 7.13	69	-12.2	-16.13	-8.48	<0.001	99.419
ADF	310.9 ± 6.41	305.2 ± 5.94	70	-5.7	-8.64	-3.01	<0.001	98.830
NH ₃ -N (g/kg total N)	130.3 ± 5.75	84.8 ± 4.97	70	-45.5	-55.50	-34.74	<0.001	99.764
CP	223.6 ± 3.50	227.8 ± 3.39	53	4.2	2.83	6.19	<0.001	98.422
WSC	14.7 ± 1.36	13.3 ± 1.32	67	-0.31	-0.81	0.20	0.240	98.769
Ash	134.8 ± 1.75	133.1 ± 1.63	13	-1.4	-2.90	0.67	0.221	95.695
IVDMD-48 h	669.2 ± 9.55	675.3 ± 9.55	11	6.1	-9.13	20.29	0.457	93.858
Ethanol	5.8 ± 0.51	3.5 ± 0.53	28	-2.3	-3.02	-1.68	<0.001	99.855
Lactate	39 ± 3.36	46.5 ± 3.21	119	7.5	5.82	8.07	<0.001	99.230
Acetate	20.4 ± 1.37	19.3 ± 1.48	94	-1.1	-2.27	-1.22	<0.001	99.544
Propionate	3.5 ± 0.79	2.7 ± 0.76	44	-0.8	-1.55	-0.80	<0.001	99.804
Butyrate	7.5 ± 0.81	2.7 ± 0.78	29	-4.8	-4.28	-2.92	<0.001	99.291
LAB (log ₁₀ cfu/g)	7.1 ± 1.32	7 ± 1.30	37	-0.1	-0.36	0.02	0.077	99.945
Yeast (log ₁₀ cfu/g)	3.1 ± 0.31	2.5 ± 0.21	14	-0.6	-1.12	-0.13	0.013	95.335
Mould (log ₁₀ cfu/g)	4.1 ± 0.32	2.2 ± 0.32	12	-1.9	-3.35	-0.75	0.002	99.721
Aerobic stability (h)	237 ± 20.10	263.7 ± 20.10	10	26.7	13.42	39.98	<0.001	41.307

RMD, raw mean difference between inoculated and uninoculated treatments; DM, dry matter; NDF, neutral detergent fibre; ADF, acid detergent fibre; NH₃-N, ammoniacal nitrogen; CP, crude protein; WSC, water-soluble carbohydrates; IVDMD-48 h, *in vitro* DM digestibility at 48 h; LAB, lactic acid bacteria.

Results were expressed as the arithmetic mean ± standard error of the mean (S.E.M). Bold values denote statistical significance at the $P < 0.05$ level.

the absence of *P. acidilactici*. Silage inoculation with *P. acidilactici* tended to increase DM content ($P = 0.092$). Meanwhile, *P. acidilactici* inoculation reduced pH, NDF, ADF, butyrate and yeast counts. CP and lactate concentrations increased, whereas ethanol decreased ($P < 0.05$) both in the presence and in the absence of *P. acidilactici* in the inoculated group (Table 2).

Inoculation with *E. faecium* significantly reduced WSC ($P = 0.017$), whereas its absence decreased ADF, acetate, butyrate, yeasts, moulds ($P < 0.05$) and tended to decrease propionate concentrations ($P = 0.061$). On the contrary, DM and CP increased, while ethanol decreased in the absence of this microorganism ($P < 0.001$). pH, NDF, NH₃-N were reduced ($P < 0.05$), while lactate was increased ($P < 0.001$) both in the presence and in the absence of *E. faecium* in the inoculated group (Table 2).

Based on the results from the meta-regression analysis, interactions were observed between the year of publication and CP ($P < 0.001$), WSC ($P = 0.001$), acetate ($P = 0.002$), propionate ($P = 0.001$) and yeasts ($P = 0.004$). Moreover, the application rate of LAB was associated with pH ($P = 0.001$), DM ($P = 0.043$), CP ($P = 0.014$), WSC ($P = 0.016$), ash ($P = 0.021$) and propionate ($P = 0.015$). Finally, the duration of studies was correlated with pH ($P < 0.001$), DM ($P = 0.0002$), NH₃-N ($P < 0.001$), WSC ($P < 0.001$), ash ($P = 0.001$), ethanol ($P = 0.024$), acetate ($P < 0.001$), propionate ($P = 0.0003$), butyrate ($P = 0.005$) and yeast counts ($P = 0.018$) in the meta-regression (Table 4). Yet, the coefficient was under 50%, except for the duration of studies in WSC (adjusted $R^2 = 0.72$). Therefore, the year, rate of inoculation and length of studies had a reduced impact in the remaining response variables.

As part of this study, beside Egger's regression test, Begg and Mazumdar rank correlation test, Duval and Tweedie's trim and fill method were used to detect publication bias in the studies included for each of the response variables analysed. The results are shown in Table 5. There was a general tendency in having few publication biases for pH, CP and butyrate; however, the large number of scientific articles included in this meta-analysis provides valid results beyond the potential bias (Table 5).

Discussion

This quantitative meta-analysis of data from several randomized controlled experiments showed that the use of LAB inoculants decreased pH in the pooled estimate. Results also suggested that the nutrients in alfalfa silage were well preserved by inoculation with LAB, as indicated by lower NDF and ADF concentrations, and higher DM and CP in the pooled estimate as compared with the control silage. The NDF and ADF of silages are important quality parameters and are expected to be lower in inoculated alfalfa. Certain inoculants contain bacteria that could secrete specific enzymes, mainly cellulases and xylanases, that may contribute to degrade these structures and increased fibre digestion (Adesogan *et al.*, 2019). The decreased NDF and ADF concentration could be also related to better WSC preservation following inoculation, which could reduce NDF and ADF contents by a 'dilution effect'. The reduction of NDF and ADF in treated silage compared with the control evidenced favourable anaerobic conditions for the fermentation process, degradation of cell walls providing soluble carbohydrates to fermentative microorganisms

Table 2. Sub-group analysis comparing the effects of silage inoculation on nutrient composition and microbiological profile of alfalfa silage according to the LAB species

Response variables	HoLAB	HeLAB	With <i>L. buchneri</i>	Without <i>L. buchneri</i>	With <i>L. plantarum</i>	Without <i>L. plantarum</i>	With <i>P. acidilactici</i>	Without <i>P. acidilactici</i>	With <i>E. faecium</i>	Without <i>E. faecium</i>
pH	-0.4 ± 0.04 (43)	-0.2 ± 0.12 (8)	-0.1 ± 0.07 (17)	-0.3 ± 0.03 (70)	-0.2 ± 0.03 (61)	-0.4 ± 0.07 (26)	-0.2 ± 0.06 (15)	-0.3 ± 0.03 (72)	-0.2 ± 0.04 (13)	-0.3 ± 0.03 (74)
DM	8.2 ± 1.45 (40)	-4.4 ± 4.57 (8)	-5.0 ± 2.85 (13)	8.0 ± 1.11 (71)	6.5 ± 1.07 (61)	4.3 ± 2.56 (23)	6.0 ± 3.53 (23)	6.0 ± 1.02 (61)	-1.1 ± 1.62 (17)	7.8 ± 1.27 (67)
NDF	-15.7 ± 2.52 (32)	-29.9 ± 30.00 (2)	-9.7 ± 5.93 (8)	-13.2 ± 2.15 (58)	-8.8 ± 2.52 (50)	-23.2 ± 3.53 (16)	-20.8 ± 5.63 (18)	-10.2 ± 2.65 (48)	-8.4 ± 4.01 (19)	-14.2 ± 2.14 (47)
ADF	-7.5 ± 2.26 (34)	-17.1 ± 40.05 (2)	-5.6 ± 9.62 (6)	-6.5 ± 1.51 (61)	-3.5 ± 1.60 (53)	-15.5 ± 3.39 (14)	-13.4 ± 3.86 (19)	-4.2 ± 1.81 (48)	-0.3 ± 2.11 (19)	-8.1 ± 1.90 (48)
NH ₃ -N (g/kg total N)	-45.3 ± 5.85 (38)	-31.8 ± 13.28 (5)	-27.6 ± 11.43 (6)	-33.4 ± 4.08 (55)	-25.5 ± 4.06 (41)	-47.7 ± 8.28 (20)	-21.6 ± 14.27 (9)	-34.6 ± 3.91 (52)	-7.7 ± 4.10 (12)	-39.2 ± 4.61 (49)
CP	5.3 ± 1.03 (30)	14.8 ± 9.00 (2)	5.2 ± 2.25 (7)	4.3 ± 0.96 (42)	2.4 ± 1.79 (34)	7.9 ± 1.15 (15)	12.6 ± 4.68 (6)	3.4 ± 1.03 (43)	3.5 ± 3.86 (8)	4.5 ± 0.98 (41)
WSC	-0.6 ± 0.37 (39)	1.0 ± 0.20 (8)	0.7 ± 0.23 (14)	-0.7 ± 0.37 (49)	-0.5 ± 0.34 (46)	0.3 ± 0.44 (17)	-1.8 ± 1.26 (10)	-0.1 ± 0.28 (53)	-4.0 ± 1.68 (3)	-0.1 ± 0.27 (60)
Ash	-2.3 ± 1.44 (5)	-0.5 ± 1.45 (8)	2.5 ± 1.32 (4)	-2.9 ± 1.10 (9)	-1.5 ± 1.27 (11)	0.0 ± 2 (2)	-1.1 ± 2.39 (3)	-1.2 ± 1.07 (10)	-3.8 ± 2.76 (3)	-0.6 ± 1.12 (10)
IVDMD-48 h	5.6 ± 7.51 (11)	-	-11.0 ± 4.60 (4)	14.7 ± 11.63 (6)	1.1 ± 8.33 (9)	36.0 ± 8.49 (1)	0.2 ± 42.50 (2)	5.9 ± 6.45 (8)	-	-0.6 ± 7.88 (9)
Ethanol	-2.1 ± 0.92 (10)	-1.8 ± 1.41 (6)	-0.9 ± 0.40 (11)	-3.3 ± 0.64 (17)	-2.8 ± 0.38 (21)	-0.7 ± 1.41 (7)	-3.1 ± 0.50 (1)	-2.3 ± 0.35 (27)	-2.5 ± 2.19 (3)	-2.3 ± 0.43 (25)
Lactate	4.9 ± 0.82 (53)	3.8 ± 1.09 (9)	0.6 ± 0.98 (16)	6.7 ± 0.68 (84)	5.3 ± 0.56 (75)	5.8 ± 1.83 (25)	8.8 ± 2.02 (23)	4.9 ± 0.54 (77)	9.2 ± 1.92 (19)	4.8 ± 0.54 (81)
Acetate	-2.5 ± 0.41 (46)	10.8 ± 2.84 (7)	4.2 ± 0.44 (15)	-2.8 ± 0.34 (77)	-2.9 ± 0.32 (62)	0.9 ± 0.72 (30)	-1.7 ± 1.17 (15)	-1.8 ± 0.29 (77)	0.0 ± 1.22 (13)	-2.0 ± 0.29 (79)
Propionate	-0.5 ± 0.61 (17)	-0.8 ± 1.78 (5)	-0.1 ± 1.05 (11)	-0.6 ± 0.22 (24)	-0.9 ± 0.16 (22)	-0.1 ± 1.16 (13)	1.1 ± 2.66 (6)	-0.8 ± 0.25 (29)	0.1 ± 0.25 (3)	-0.5 ± 0.28 (32)
Butyrate	-5.5 ± 1.13 (14)	-11.3 ± 6.49 (3)	-2.2 ± 0.36 (9)	-4.7 ± 0.67 (17)	-1.5 ± 0.37 (19)	-9.3 ± 1.42 (7)	-12.3 ± 7.36 (4)	-2.5 ± 0.35 (22)	0.02 ± 0.07 (1)	-3.9 ± 0.40 (25)
LAB (log ₁₀ cfu/g)	-0.4 ± 0.34 (14)	-0.002 ± 0.29 (3)	0.3 ± 0.07 (8)	-0.4 ± 0.23 (26)	-0.1 ± 0.06 (24)	-0.3 ± 0.45 (10)	-1.2 ± 1.46 (2)	-0.1 ± 0.05 (32)	-0.2 ± 0.15 (5)	-0.2 ± 0.11 (29)
Yeast (log ₁₀ cfu/g)	-1.0 ± 0.29 (5)	-1.0 ± 0.23 (3)	-1.3 ± 0.23 (4)	-0.4 ± 0.30 (10)	-0.5 ± 0.31 (11)	-1.0 ± 0.23 (3)	-0.9 ± 0.21 (1)	-0.6 ± 0.27 (13)	0.5 ± 0.28 (1)	-0.7 ± 0.26 (13)
Mould (log ₁₀ cfu/g)	-3.4 ± 0.21 (4)	-2.9 ± 1.16 (2)	-2.8 ± 0.46 (4)	-1.7 ± 0.85 (8)	-1.9 ± 0.73 (10)	-2.9 ± 1.14 (2)	-	-2.1 ± 0.66 (12)	-0.1 ± 0.32 (2)	-2.5 ± 0.28 (10)

RMD, raw mean difference between inoculated and uninoculated treatments; HoLAB, homofermentative lactic acid bacteria; HeLAB, heterofermentative lactic acid bacteria; DM, dry matter; NDF, neutral detergent fibre; ADF, acid detergent fibre; NH₃-N, ammoniacal nitrogen; CP, crude protein; WSC, water-soluble carbohydrates; IVDMD-48 h, *in vitro* DM digestibility at 48 h; LAB, lactic acid bacteria.

Results were expressed as the RMD ± S.E.M and values in parentheses indicate the number of studies. There was no sub-group analysis for aerobic stability.

Table 3. Sub-group analysis comparing the effects of silage inoculation on nutrient composition and microbiological profile of alfalfa silage considering factors that could potentially influence the magnitude of the treatment

Response variables	Before 2009	After 2009	Mono-strain	Multi-strain	With enzymes	Without enzymes	30 to 60 days	60 days	Mini silos	Farm scale
pH	-0.4 ± 0.05 (41)	-0.3 ± 0.03 (65)	-0.4 ± 0.04 (49)	-0.3 ± 0.04 (57)	0.0 ± 0.07 (7)	-0.4 ± 0.03 (99)	-0.4 ± 0.04 (68)	-0.3 ± 0.05 (35)	-0.4 ± 0.03 (96)	-0.2 ± 0.05 (10)
DM	6.4 ± 2.42 (38)	6.0 ± 1.11 (50)	5.5 ± 1.35 (45)	6.6 ± 1.63 (42)	11.5 ± 4.18 (14)	5.5 ± 1.01 (74)	9.3 ± 1.42 (46)	2.1 ± 1.39 (39)	5.0 ± 0.96 (78)	17.4 ± 12.85 (10)
NDF	-10.0 ± 4.36 (25)	-13.3 ± 2.33 (44)	-19.2 ± 2.53 (31)	-5.9 ± 2.71 (38)	-13.7 ± 4.99 (18)	-12.0 ± 2.05 (51)	-8.4 ± 2.49 (34)	-17.8 ± 5.80 (33)	-11.4 ± 2.09 (60)	-18.4 ± 7.72 (9)
ADF	-5.0 ± 2.26 (27)	-6.2 ± 1.76 (43)	-10.6 ± 2.23 (33)	-0.6 ± 2.09 (37)	-5.5 ± 2.53 (16)	-5.7 ± 1.82 (54)	-3.1 ± 1.78 (35)	-8.5 ± 5.23 (34)	-6.0 ± 1.55 (60)	-4.8 ± 3.71 (10)
NH ₃ -N (g/kg total N)	-55.8 ± 14.21 (28)	-38.4 ± 4.71 (42)	-43.9 ± 5.64 (39)	-40.6 ± 4.80 (30)	-12.5 ± 11.72 (4)	-47.1 ± 5.65 (66)	-53.3 ± 7.46 (44)	-31.4 ± 6.61 (26)	-50.3 ± 5.80 (61)	-10.8 ± 5.30 (9)
CP	-1.3 ± 2.31 (9)	5.5 ± 0.93 (44)	5.7 ± 1.05 (29)	2.1 ± 2.39 (23)	0.7 ± 4.57 (8)	5.1 ± 0.86 (45)	3.0 ± 0.92 (32)	7.6 ± 3.54 (19)	5.6 ± 0.85 (48)	-6.7 ± 4.24 (5)
WSC	-2.8 ± 0.83 (25)	0.1 ± 0.29 (42)	-0.5 ± 0.35 (39)	0.0 ± 0.41 (27)	-0.2 ± 0.29 (4)	-0.3 ± 0.27 (63)	-0.6 ± 0.22 (39)	0.8 ± 0.25 (28)	0.0 ± 0.26 (60)	-7.6 ± 2.51 (7)
Ash	-	-1.1 ± 0.91 (13)	-2.3 ± 1.44 (5)	-0.5 ± 1.45 (8)	-0.1 ± 1.73 (6)	-2.1 ± 1.27 (7)	0.3 ± 1.09 (4)	-1.7 ± 1.90 (8)	-0.7 ± 0.89 (12)	-5.0 ± 0.67 (1)
IVDMD-48 h	16.4 ± 18.99 (2)	3.2 ± 9.12 (9)	-	5.6 ± 7.51 (11)	-12.4 ± 11.58 (5)	21.1 ± 10.97 (6)	13.0 ± 17.28 (5)	-6.1 ± 6.29 (5)	2.3 ± 10.12 (9)	20.3 ± 14.45 (2)
Ethanol	-3.4 ± 1.57 (8)	-1.9 ± 0.40 (20)	-1.8 ± 0.74 (15)	-2.8 ± 0.30 (13)	-1.1 ± 1.07 (4)	-2.6 ± 0.38 (24)	-2.4 ± 0.97 (12)	-2.6 ± 0.52 (14)	-2.4 ± 0.39 (27)	-0.3 ± 0.04 (1)
Lactate	8.1 ± 1.58 (52)	6.4 ± 0.50 (67)	6.0 ± 0.70 (53)	7.4 ± 1.22 (66)	8.6 ± 1.88 (17)	6.6 ± 0.61 (100)	6.4 ± 1.46 (77)	7.6 ± 0.55 (39)	6.3 ± 0.60 (106)	12.6 ± 1.73 (13)
Acetate	-1.3 ± 0.45 (40)	-2.1 ± 0.29 (54)	-1.0 ± 0.38 (50)	-2.4 ± 0.56 (44)	-4.5 ± 1.92 (13)	-1.2 ± 0.27 (81)	-1.2 ± 0.37 (58)	-2.7 ± 0.34 (33)	-1.1 ± 0.27 (84)	-7.3 ± 2.64 (10)
Propionate	-1.8 ± 0.34 (21)	-0.5 ± 0.32 (23)	-0.7 ± 0.76 (19)	-1.6 ± 0.17 (24)	0.0 ± 0.01 (4)	-1.3 ± 0.28 (40)	-1.6 ± 0.32 (24)	-0.6 ± 0.40 (19)	-1.3 ± 0.26 (40)	-0.1 ± 0.14 (4)
Butyrate	-1.0 ± 0.25 (10)	-4.9 ± 0.58 (19)	-9.5 ± 1.33 (13)	0.0 ± 0.28 (16)	0.1 ± 0.05 (3)	-4.4 ± 0.44 (26)	-0.3 ± 1.24 (9)	-5.7 ± 0.43 (19)	-4.5 ± 0.48 (25)	-0.3 ± 0.18 (4)
LAB (log ₁₀ cfu/g)	-0.3 ± 0.08 (15)	-0.1 ± 0.13 (22)	-0.3 ± 0.27 (17)	-0.0 ± 0.04 (20)	0.4 ± 0.03 (4)	-0.2 ± 0.20 (33)	-0.2 ± 0.08 (18)	-0.2 ± 0.13 (19)	-0.1 ± 0.10 (35)	-0.9 ± 0.25 (2)
Yeast (log ₁₀ cfu/g)	0.6 ± 0.10 (3)	-0.1 ± 0.19 (11)	-1.0 ± 0.19 (8)	-0.1 ± 0.41 (6)	0.5 ± 0.28 (1)	-0.7 ± 0.26 (13)	0.6 ± 0.10 (3)	-1.0 ± 0.19 (11)	-0.6 ± 0.25 (14)	-
Mould (log ₁₀ cfu/g)	-1.3 ± 0.19 (3)	-2.3 ± 0.79 (9)	-3.1 ± 0.31 (5)	-1.4 ± 0.79 (7)	-0.6 ± 0.35 (1)	-2.2 ± 0.70 (11)	-1.3 ± 0.19 (3)	-2.3 ± 0.79 (9)	-2.1 ± 0.66 (12)	-

RMD, raw mean difference between inoculated and uninoculated treatments; DM, dry matter; NDF, neutral detergent fibre; ADF, acid detergent fibre; NH₃-N, ammonia nitrogen; CP, crude protein; WSC, water-soluble carbohydrates; IVDMD-48 h, *in vitro* DM digestibility at 48 h; LAB, lactic acid bacteria.

Results were expressed as the RMD ± S.E.M and values in parentheses indicate the number of studies. There was no sub-group analysis for aerobic stability.

Table 4. Summary of random weighted meta-regression analysis for independent variables (year of publication, application rate of LAB and duration of studies) that influenced the effects between inoculated and uninoculated treatments for alfalfa silage quality parameters (g/kg DM, unless otherwise stated)

Response variable	Meta-regression parameters											
	Year of publication				Application rate of LAB				Duration of studies			
	Intercept ^a	Slope	<i>P</i> value	<i>I</i> ² (%)	Intercept ^a	Slope	<i>P</i> value	<i>I</i> ² (%)	Intercept ^a	Slope	<i>P</i> value	<i>I</i> ² (%)
pH	2.2615	-0.0013	0.622	99.5	-1.1823	0.1539	0.001	99.56	-0.6374	0.0047	<0.001	99.46
DM	-148.19	0.0767	0.389	96	5.47	<0.0001	0.043	96.06	2.302	-0.022	0.0002	94.68
NDF	125.45	-0.069	0.726	99.43	0.7118	<-0.0001	0.712	99.4	-10.051	-0.0346	0.525	99.42
ADF	188.779	-0.0097	0.505	98.85	-4.8099	<0.0001	0.079	98.89	-9.9745	0.0607	0.170	98.86
NH ₃ -N (g/kg total N)	-35.4664	0.0154	0.772	99.77	-5.074	<0.0001	0.058	99.77	-10.3215	0.0972	<0.001	99.7
CP	-1046.22	0.5221	<0.001	98.34	3.3248	<0.0001	0.014	97.5	3.6452	0.01559	0.563	98.28
WSC	-293.025	0.1456	<0.001	98.78	-0.166	<0.0001	0.016	98.41	-1.2705	0.02192	<0.001	95.46
Ash	-804.529	0.3988	0.429	95.66	0.64506	<-0.001	0.021	95.11	11.2058	-0.1654	0.001	94.67
IVDMD-48 h	2394.895	-1.19003	0.128	93.78	-8.5457	0.00003	0.251	95.13	51.714	-0.8125	0.111	93.94
Ethanol	-26.6216	0.01208	0.823	99.83	-2.5656	<0.0001	0.378	99.79	1.6928	-0.0596	0.024	99.86
Lactate	189.687	-0.0910	0.066	98.84	6.5925	<-0.0001	0.076	99.07	7.0987	-0.0026	0.889	99.12
Acetate	150.414	-0.0758	0.002	99.41	1.383	<0.0001	0.175	99.56	3.1292	-0.0805	<0.001	99.24
Propionate	-157.363	0.0778	0.001	99.81	-1.597	<0.0001	0.015	99.83	-3.7992	0.0440	0.0003	99.8
Butyrate	82.4944	-0.0429	0.130	99.31	-3.5147	<-0.0001	0.201	99.4	-5.5241	0.0326	0.005	99.34
LAB (log ₁₀ cfu/g)	12.4227	-0.0063	0.595	99.94	0.3525	-0.0975	0.345	99.92	-0.2285	0.0008	0.763	99.94
Yeast (log ₁₀ cfu/g)	244.8756	-0.122	0.004	91.13	-1.8489	0.1994	0.748	95.84	1.9948	-0.0336	0.018	92.16
Mould (log ₁₀ cfu/g)	211.2289	-0.106	0.556	99.72	-3.651	0.2625	0.844	99.64	1.3897	-0.0419	0.352	99.61

LAB, lactic acid bacteria; DM, dry matter; NDF, neutral detergent fibre; ADF, acid detergent fibre; NH₃-N, ammoniacal nitrogen; CP, crude protein; WSC, water-soluble carbohydrates; IVDMD-48 h, *in vitro* DM digestibility at 48 h. Bold values denote statistical significance at the *P* < 0.05 level.

^aIntercept: constant in the model.

Table 5. Publication bias detection (g/kg DM, unless otherwise stated)

Response variable	Fail-safe N ^a	Begg and Mazumdar test	Egger's regression test	
			Intercept	P value
pH	0	0.01629	-7.21081	0.001
DM	1	0.52587	0.97517	0.201
NDF	3	0.65226	-1.71318	0.361
ADF	5	0.18243	-0.07919	0.952
NH ₃ -N (g/kg total N)	0	0.06953	-12.01304	0.003
CP	0	0.00422	-3.91795	0.002
WSC	0	0.9224	-2.60699	0.060
Ash	0	0.36012	-1.97912	0.415
IVDMD-48 h	0	0.31151	0.39048	0.873
Ethanol	0	0.64954	-14.38607	0.005
Lactate	0	0.26629	-0.71528	0.544
Acetate	4	<0.001	2.9993	0.139
Propionate	0	0.37344	-6.49239	0.089
Butyrate	0	0.03733	-7.0714	0.021
LAB (log ₁₀ cfu/g)	0	0.00579	-14.29756	0.077
Yeast (log ₁₀ cfu/g)	0	0.82667	-9.76489	0.302
Mould (log ₁₀ cfu/g)	0	0.1314	-11.04397	0.186

DM, dry matter; NDF, neutral detergent fibre; ADF, acid detergent fibre; NH₃-N, ammoniacal nitrogen; CP, crude protein; WSC, water-soluble carbohydrates; IVDMD-48 h, *in vitro* DM digestibility at 48 h; LAB, lactic acid bacteria.

^aNumber of studies required to reverse the effects are calculated on the condition of $P = 0.05$.

and hydrolysis of the most available forage structural carbohydrates (Kozelov *et al.*, 2008; Rabelo *et al.*, 2016; Wang *et al.*, 2019). Bearing in mind the sub-group analysis, although some categories did not show significant differences, in all cases, there was a reduction in both parameters.

Independently of the factors that could influence the treatment magnitude, DM content increased, except when experiments lasted for more than 60 days and when HeLAB were applied. The lack of effect with prolonged storage time can be related to the degradation of some nutrients and cell walls by bacterial enzymes activity and acidic conditions in silage during fermentation (Sarıççek *et al.*, 2016). On the other hand, since obligate HeLAB are characterized by CO₂ production in the conversion of lactic acid and carbohydrates to acetic and propionic acids, they would decrease DM content (Weinberg *et al.*, 2010; Randby *et al.*, 2012). The higher CP and the lower content of NH₃-N indicated limited proteolysis. In this meta-analysis, LAB inoculants exhibited their potential to protect feed proteins in alfalfa silage. The positive effects of LAB inoculants on nitrogen fractions can be accounted for the rapid acidification of the forage below the optimal pH for plant protease activity (Wang *et al.*, 2009). During ensiling, plant enzymes such as carboxypeptidase (optimum pH 5.2) and acid proteinase (optimum pH 4.5) may play major roles in protein degradation, while aminopeptidase loses most of its activity in the initial phase of fermentation since its optimal pH is close to 7.0 (McKersie and

Buchanan-Smith, 1982). Accordingly, if the pH of forage is reduced as rapidly as possible, it will contribute to proteolysis inhibition (Wang *et al.*, 2009).

With reference to organic acids, in treated silages, lactic acid concentrations were largely increased, whereas the levels of ethanol, acetic, propionic and butyric acids were clearly lower than in uninoculated silages. LAB have a positive effect on the extent and rate of lactic acid production in the silage, hence stimulating a rapid drop in pH and suppressing the growth of clostridia and other undesired anaerobic microorganisms (Oude Elferink *et al.*, 2001). Although inoculation with LAB reduced acetate concentrations, overall HeLAB significantly increased it. While inoculation with HoLAB led to silages with high lactic acid contents, inoculation with HeLAB resulted in higher levels of acetic acid (Chen *et al.*, 2018). It is well documented (Heinl and Grabherr, 2017) that this could be due to the capacity of HeLAB to degrade lactic acid to acetic acid under anoxic conditions. Acetic acid is one of the most effective substances for inhibition of spoilage microorganisms by decreasing their maximum growth rate (Danner *et al.*, 2003). A level of acetic acid of 1.5–3.0% in the DM could inhibit yeast growth in silages exposed to air in the feed out phase (Acosta Aragón *et al.*, 2012). In the same way, propionic acid levels were lower in inoculated silages. Although a certain amount of propionic acid is desirable in order to minimize possible growth of yeasts and improve aerobic stability, it could affect the voluntary intake and utilization of silage-based diets (Nishino *et al.*, 2003). In conserved forage, butyric acid and ethanol are equally undesirable. In the present work, lower amounts of ethanol appeared in inoculated alfalfa silages. Ethanol, which is a yeast end product, has little preservative effect in silage, and it causes extremely high losses in DM and energy (Kung *et al.*, 2018). With respect to butyrate, this organic acid confers poor palatability, reducing the voluntary feed intake in animals (Kung, 2010). Frequently, uninoculated silages have relatively high contents of butyrate, related to the activity of clostridia derived from soil or slurry contamination (Danner *et al.*, 2003; Liu *et al.*, 2016).

Taking into account the microbiological composition, all treated silages had lower counts of yeasts and moulds than the control ones in the pool estimate. On the contrary, the inclusion of inoculants did not significantly raise LAB counts. However, it should be noted that LAB reached a concentration of at least 10⁶ cfu/g in either group after ensiling. It was also worth noting that, regardless of the number of final LAB, when an inoculant fails to produce an adequate amount of lactic acid in the process of silage fermentation to reduce pH and suppress the growth of harmful microorganisms, the resulting silage will be of poor quality (Ni *et al.*, 2015). In the studies summarized in this analysis, the counts of LAB were similar, but it was possible to detect significant differences in the aforementioned variables in favour of the treated group. Moreover, it was possible that differences in the numbers of LAB were only seen in the early stages of fermentation. The fast increase in LAB counts frequently observed in inoculated silages in early fermentation indicates that LAB strains are competitive among the epiphytic communities. Moreover, the reduction in the LAB population after this initial peak is expected because low pH and lack of fermentable substrates result in bacterial death (Xu *et al.*, 2017; Nascimento Agarussi *et al.*, 2019).

Regarding yeasts, these microorganisms are the main initiators of aerobic spoilage by metabolizing valuable sugars and lactic acid, thus raising the pH and allowing an increase of silage inner temperature (Pahlow *et al.*, 2003). Finally, moulds complete the deterioration of silages (Dolci *et al.*, 2011). For well-preserved

silage, the concentration of moulds and yeasts should not exceed 3–4 log₁₀ cfu/g (McEniry *et al.*, 2006). In the present study, inoculating alfalfa at the time of ensiling altered the resulting fermentation by reducing the level of yeasts and moulds in 1–2 log₁₀ cfu/g compared with untreated silage. On the other hand, yeast counts were increased in studies conducted for less than 60 days. Apparently, the slow development of HeLAB, which are the main antifungal agents, explains why the effects of these microorganisms are manifest only during the late storage phase of ensiling (Schmidt *et al.*, 2009).

The results observed in the pool estimate have led us to hypothesize that alfalfa silage inoculants provide a stable acidic pH with a suitable proportion of organic acids after opening the fermented plant material. Thereby, growth of yeasts and moulds in the presence of oxygen is inhibited and also heating of the silage is prevented. Classical microbial inoculants containing only HoLAB were shown to have no significant influence on aerobic stability, primarily because lactic acid by itself is not an effective antimycotic agent (Filya and Sucu, 2007). A more promising approach seems to be related to the use of HeLAB. Addition of these microorganisms improves aerobic stability through the production of acetic and propionic acids with strong antifungal properties (Zielińska *et al.*, 2015). However, our meta-analysis summarized limited reports on the ability of microbial inoculants to improve the aerobic stability of alfalfa silage, and it was not possible to perform a sub-group analysis to investigate its influence on the response. Hence, future research should be conducted to further examine the effects of HoLAB, HeLAB and their combinations during aerobic exposure. Although the inclusion of HeLAB on forages with a low DM content does not appear to be appropriate due to the excessive fermentation (Jatkauskas *et al.*, 2013), the aforementioned inoculants should be utilized so as to avoid the air deterioration that could occur during the feed out phase or due to poor management (Yuan *et al.*, 2018). For instance, silage moved from one silo structure to another, silage fed from intermediate feeding piles, and silos with large exposed surfaces are good candidates for treatment with HeLAB. Besides, over-sized silos, with slow feed out rate, poor packing and maintained at 30°C, are more prone to aerobic deterioration, so combinations of several strains with different mechanisms of action should be considered (Ashbell *et al.*, 2002; Kung, 2010).

According to the sub-group analyses, the use of enzymes did not offer supplemental effects on silage pH. The result in studies applying enzymes could be attributed to the fact that all the trials involved employed HeLAB, too. It might be hypothesized that the HeLAB *L. buchneri* was a confounder and had possibly a major influence on this effect, as using this microorganism implies a more heterolactic fermentation, consisting in the conversion of pentoses or hexoses into lactic acid, CO₂ and other products, mainly acetic acid, ethanol and propionic acid (McDonald *et al.*, 1991). Furthermore, the absence of enzymes significantly increased CP and IVDMD-48 h, while decreased NH₃-N, ethanol, propionate, yeasts and moulds counts. Therefore, this meta-analysis indicates no benefit of the LAB treatment with combined fibrolytic enzymes. Though this is in agreement with the results obtained by Lynch *et al.* (2015), one plausible explanation could be that the number of experiments that employed enzymes and measured certain variables was relatively small, thus this observation should be interpreted with caution.

Summary of findings included in this meta-analysis denoted that *L. buchneri*, *L. plantarum*, *P. acidilactici* and *E. faecium* were mostly administered as multi-species inocula due to the

synergistic effects when bacteria are applied together (Blajman *et al.*, 2018). Therefore, the infrequent use of individual inoculants (except for *L. plantarum*) may have limited our ability to detect LAB species-related impacts on the measures of silage quality. Still, this meta-analysis evidenced that the main goal of *L. plantarum*, *P. acidilactici* and *E. faecium* administration was the preservation of the nutritional quality of ensiled alfalfa (Oliveira *et al.*, 2017), whereas reduction in harmful microorganisms was the most consistent benefit of *L. buchneri* (Liu *et al.*, 2018).

This work produced a synthesis and contrast of results among a large number of primary studies. To our knowledge, this is the first meta-analysis to compare the addition of HoLAB and HeLAB for alfalfa silage. In the pool estimate, positive effects due to the application of microbial silage inoculants were found in most of the evaluated parameters. Regarding the sub-group analysis, inoculation with HoLAB is recommended as it has been shown to contribute to a lesser loss of nutritional value and to improve the chemical parameters of alfalfa silages. Moreover, this meta-analysis provided evidence that using either HoLAB or HeLAB enhanced microbiological composition. In spite of the previous statement, further studies are needed to examine the effects of HoLAB and HeLAB with different biotechnological features and in appropriate proportions on digestibility and animal performance. Additionally, more studies are required to identify the effects of LAB inoculants on silage preserved in farm-scale silos as well as on how LAB combined with enzymes affects silage quality. Lastly, research should be conducted to analyse the ability of bio-inoculants to inhibit clostridia and decontaminate silages of mycotoxins produced by them, so as to finally standardize commercial alfalfa silage inoculants.

Acknowledgements. The authors would like to thank Lecturer Liliana Silber for her English manuscript proofreading service.

Financial support. This research was financially supported by the National Institute of Agricultural Technology (INTA, Argentina) and The National Agency of Scientific and Technological Promotion (FONCyT, Argentina), Projects PICT-2017-0665 and PICT-2016-0256.

Conflict of interest. None.

Ethical standards. Not applicable.

References

- Acosta Aragón Y, Jatkauskas J and Vrotniakienė V (2012) The effect of a silage inoculant on silage quality, aerobic stability, and meat production on farm scale. *International Scholarly Research Notices: Veterinary Science* **2012**, 1–6.
- Adesogan AT, Arriola KG, Jiang Y, Oyebade A, Paula EM, Pech-Cervantes AA, Romero JJ, Ferraretto LF and Vyas D (2019) Symposium review: technologies for improving fibre utilization. *Journal of Dairy Science* **102**, 5726–5755.
- Arriola KG, Oliveira AS, Ma ZX, Lean IJ, Giurcanu MC and Adesogan AT (2017) A meta-analysis on the effect of dietary application of exogenous fibrolytic enzymes on the performance of dairy cows. *Journal of Dairy Science* **100**, 1–15.
- Ashbell G, Weinberg ZG, Hen Y and Filya I (2002) The effects of temperature on the aerobic stability of wheat and corn silages. *The Journal of Industrial Microbiology and Biotechnology* **28**, 261–263.
- Begg CB and Mazumdar M (1994) Operating characteristics of rank correlation test for publication bias. *Biometrics* **50**, 1088–1101.
- Blajman JE, Páez RB, Vinderola CG, Lingua MS and Signorini ML (2018) A meta-analysis on the effectiveness of lactic acid bacteria for corn silage. *Journal of Applied Microbiology* **125**, 1655–1669.

- Borenstein M, Hedges LV, Higgins JPT and Rothstein HR** (2009) *Introduction to Meta-Analysis*. Chichester, West Sussex, UK: John Wiley & Sons, Ltd.
- Borreani G, Tabacco E, Schmidt RJ, Holmes BJ and Muck RE** (2018) Silage review: factors affecting dry matter and quality losses in silages. *Journal of Dairy Science* **101**, 3952–3979.
- Chen L, Dong Z, Li J and Shao T** (2018) Ensiling characteristics, *in Vitro* rumen fermentation, microbial communities and aerobic stability of low-dry matter silages produced with sweet sorghum and alfalfa mixtures. *Journal of the Science of Food and Agriculture* **99**, 2140–2151.
- Chilson JM, Rezam P, Drewnoski M, Price W and Hunt CW** (2016) Effect of homofermentative lactic acid bacteria and exogenous hydrolytic enzymes on the ensiling characteristics and rumen degradability of alfalfa and corn silages. *The Professional Animal Scientist* **32**, 598–604.
- Contreras-Govea FE, Muck RE, Mertens DR and Weimer PJ** (2011) Microbial inoculant effects on silage and *in vitro* ruminal fermentation, and microbial biomass estimation for alfalfa, bmr corn, and corn silages. *Animal Feed Science and Technology* **163**, 2–10.
- Danner H, Holzer M, Mayrhuber E and Braun R** (2003) Acetic acid increases stability of silage under aerobic conditions. *Applied and Environmental Microbiology* **69**, 562–570.
- Dolci P, Tabacco E, Cocolin L and Borreani G** (2011) Microbial dynamics during aerobic exposure of corn silage stored under oxygen barrier or polyethylene films. *Applied and Environmental Microbiology* **77**, 7499–7507.
- Dunière L, Sindou J, Chaucheyras-Durand F, Chevallier I and Thévenot-Sergentet D** (2013) Silage processing and strategies to prevent persistence of undesirable microorganisms. *Animal Feed Science and Technology* **182**, 1–15.
- Duval S and Tweedie R** (2000) Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* **56**, 455–463.
- Egger M, Davey Smith G, Schneider M and Minder C** (1997) Bias in meta-analysis detected by a simple, graphical test. *BMJ* **315**, 629–634.
- Ferrero F, Piano S, Tabacco E and Borreani G** (2019) Effects of conservation period and *Lactobacillus hilgardii* inoculum on the fermentation profile and aerobic stability of whole corn and sorghum silages. *Journal of the Science of Food and Agriculture* **99**, 2530–2540.
- Filya E and Sucu I** (2007) The effect of bacterial inoculants and a chemical preservative on the fermentation and aerobic stability of whole-crop cereal silages. *Asian-Australasian Journal of Animal Sciences* **20**, 378–384.
- Heinl S and Grabherr R** (2017) Systems biology of robustness and flexibility: *Lactobacillus buchneri* – a show case. *Journal of Biotechnology* **257**, 61–69.
- Higgins JPT and Thompson SG** (2002) Quantifying heterogeneity in a meta-analysis. *Statistics in Medicine* **21**, 1539–1558.
- Jatkaskas J, Vrotniakiene V, Ohlsson C and Lund B** (2013) The effects of three silage inoculants on aerobic stability in grass, clover-grass, lucerne and maize silages. *Agricultural and Food Science* **22**, 137–144.
- Kozelov LK, Iliev F, Hristov AN, Zaman S and McAllister TA** (2008) Effect of fibrolytic enzymes and an inoculant on *in vitro* degradability and gas production of low-dry matter alfalfa silage. *Journal of the Science of Food and Agriculture* **88**, 2568–2575.
- Kung L Jr** (2010) Aerobic stability of silage. Univ. California, Davis.
- Kung L Jr and Ranjit NK** (2001) The effect of *Lactobacillus buchneri* and other additives on the fermentation and aerobic stability of barley silage. *Journal of Dairy Science* **84**, 1149–1155.
- Kung L Jr, Shaver RD, Grant RJ and Schmidt RJ** (2018) Silage review: interpretation of chemical, microbial, and organoleptic components of silages. *Journal of Dairy Science* **101**, 4020–4033.
- Lean IJ, Rabiee AR, Duffield TF and Dohoo IR** (2009) Invited review: use of meta-analysis in animal health and reproduction: methods and applications. *Journal of Dairy Science* **92**, 3545–3565.
- Li D, Wang Y, Zhang Y, Lin Y and Yang F** (2018) Evaluation of lactic acid bacteria isolated from alfalfa for silage fermentation. *Grass Science* **64**, 190–198.
- Liu C, Lai Y, Lu X, Guo P and Luo H** (2016) Effect of lactic acid bacteria inoculants on alfalfa (*Medicago sativa* L.) silage quality: assessment of degradation (*in Situ*) and gas production (*in vitro*). *Journal of Integrative Agriculture* **15**, 2834–2841.
- Liu QH, Dong ZH and Shao T** (2018) Effect of additives on fatty acid profile of high moisture alfalfa silage during ensiling and after exposure to air. *Animal Feed Science and Technology* **236**, 29–38.
- Lynch JP, Baah J and Beauchemin KA** (2015) Conservation, fibre digestibility, and nutritive value of corn harvested at 2 cutting heights and ensiled with fibrolytic enzymes, either alone or with a ferulic acid esterase-producing inoculant. *Journal Dairy Science* **98**, 1214–1224.
- McDonald P, Henderson AR and Heron SJE** (1991) *The Biochemistry of Silage*, 2nd Edn. Aberystwyth: Cambrian Printers Ltd, pp. 184–236.
- McEniry J, O’Kiely P, Clipson NJW, Forristal PD and Doyle EM** (2006) The microbiological and chemical composition of baled and precision-chop silages on a sample of farms in County Meath. *Irish Journal of Agricultural and Food Research* **45**, 73–83.
- McKersie BD and Buchanan-Smith J** (1982) Changes in the levels of proteolytic enzymes in ensiled alfalfa forage. *Canadian Journal of Plant Science* **62**, 111–116.
- Muck RE** (2010) Silage microbiology and its control through additives. *Revista Brasileira de Zootecnia* **39**, 183–191.
- Muck RE, Nadeau EMG, McAllister TA, Contreras-Govea FE, Santos MC and Kung L Jr** (2018) Silage review: recent advances and future uses of silage additives. *Journal of Dairy Science* **101**, 3980–4000.
- Nascimento Agarussi MC, Gomes Pereira O, de Paula RA, da Silva VP, Santos Roseira JP and Fonseca e Silva F** (2019) Novel lactic acid bacteria strains as inoculants on alfalfa silage fermentation. *Scientific Reports* **9**, 8007.
- Ni K, Wang Y, Cai Y and Pang H** (2015) Natural lactic acid bacteria population and silage fermentation of whole-crop wheat. *Asian-Australasian Journal of Animal Sciences* **28**, 1123–1132.
- Nishino N, Yoshida M, Shiota H and Sakaguchi E** (2003) Accumulation of 1,2-propanediol and enhancement of aerobic stability in whole crop maize silage inoculated with *Lactobacillus buchneri*. *Journal of Applied Microbiology* **94**, 800–807.
- Ogunade IM, Kim DH, Jiang Y, Weinberg ZG, Jeong KC and Adesogan AT** (2016) Control of *Escherichia coli* O157:H7 in contaminated alfalfa silage: effects of silage additives. *Journal of Dairy Science* **99**, 4427–4436.
- Oliveira AS, Weinberg ZG, Ogunade IM, Cervantes AAP, Arriola KG, Jiang Y, Kim D, Li X, Gonçalves MCM, Vyas D and Adesogan AT** (2017) Meta-analysis of effects of inoculation with homofermentative and facultative heterofermentative lactic acid bacteria on silage fermentation, aerobic stability, and the performance of dairy cows. *Journal of Dairy Science* **100**, 4587–4603.
- Oude Elferink SJWH, Krooneman J, Gottschal JC, Spoelstra SF, Faber F and Driehuis F** (2001) Anaerobic conversion of lactic acid to acetic acid and 1,2-propanediol by *Lactobacillus buchneri*. *Applied and Environmental Microbiology* **67**, 125–132.
- Ozduven ML and Celebicam A** (2017) The effects of bacterial inoculants and/or enzymes on the fermentation characteristics and aerobic stability of alfalfa ensiled at different stages of maturity. *International Journal of Current Research* **9**, 45983–45988.
- Pahlow G, Muck RE, Driehuis F, Oude Elferink SJWH and Spoelstra SF** (2003) Microbiology of ensiling. In Buxton DR, Muck RE and Harrison HJ (eds), *Silage Science and Technology* (Agronomy Series No. 42). Madison, WI: American Society of Agronomy, pp. 31–93.
- Rabelo CHS, Mari LJ and Reis RA** (2016) Survey about the use of bacterial inoculants in Brazil: effects on silage quality and animal performance. In Silva T and Santos EM (eds), *Advances in Silage Production and Utilization*. Rijeka, Croatia: IntechOpen, pp. 3–37.
- Rabelo CHS, Lara EC, Basso FC, Härter CJ and Reis RA** (2018) Growth performance of finishing feedlot lambs fed maize silage inoculated with *Bacillus subtilis* and lactic acid bacteria. *The Journal of Agricultural Science, Cambridge* **156**, 839–847.
- Randby AT, Weisbjerg MR, Nørgaard P and Heringstad B** (2012) Early lactation feed intake and milk yield responses of dairy cows offered grass silages harvested at early maturity stages. *Journal of Dairy Science* **95**, 304–317.
- Santos MC and Kung L Jr** (2016) Short communication: the effects of dry matter and length of storage on the composition and nutritive value of alfalfa silage. *Journal of Dairy Science* **99**, 5466–5469.
- Sarıççek BZ, Yıldırım B, Kocabaş Z and Özgümüş Demir E** (2016) The effects of storage time on nutrient composition and silage quality

- parameters of corn silage made in plastic mini silo in laboratory conditions. *Iğdir University Journal of the Institute of Science and Technology* **6**, 177–183.
- Schmidt RJ, Hu W, Mills JA and Kung L Jr** (2009) The development of lactic acid bacteria and *Lactobacillus buchneri*. *Journal of Dairy Science* **92**, 5005–5010.
- Shelby LB and Vaske JJ** (2008). Understanding meta-analysis: a review of the methodological literature. *Leisure Sciences* **30**, 96–110.
- Twarużek M, Dorszewski P, Grabowicz M, Szterk P, Grajewski J and Kaszkowiak J** (2016) Effect of additives on the fermentation profile, microbial colonization and oxygen stress of alfalfa silages. *Journal of Elementology* **21**, 1161–1172.
- Wang J, Wang JQ, Zhou H and Feng T** (2009) Effects of addition of previously fermented juice prepared from alfalfa on fermentation quality and protein degradation of alfalfa silage. *Animal Feed Science and Technology* **151**, 280–290.
- Wang S, Guo G, Li J, Chen L, Dong Z and Shao T** (2019) Improvement of fermentation profile and structural carbohydrate compositions in mixed silages ensiled with fibrolytic enzymes, molasses and *Lactobacillus plantarum* MTD-1. *Italian Journal of Animal Science* **18**, 328–335.
- Weinberg ZG, Khanal P, Yildiz C, Chen Y and Arieli A** (2010) Effects of stage of maturity at harvest, wilting and LAB inoculant on aerobic stability of wheat silages. *Animal Feed Science and Technology* **158**, 29–35.
- Xu Z, He H, Zhang S and Kong J** (2017) Effects of inoculants *Lactobacillus brevis* and *Lactobacillus parafarraginis* on the fermentation characteristics and microbial communities of corn stover silage. *Scientific Reports* **7**, 13614.
- Yuan XJ, Wen AY, Wang J, Desta ST, Dong ZH and Shao T** (2018) Effects of four short-chain fatty acids or salts on fermentation characteristics and aerobic stability of alfalfa (*Medicago sativa* L.) silage. *Journal of the Science of Food and Agriculture* **98**, 328–335.
- Zhang T, Li L, Wang X, Zeng Z, Hu Y and Cui Z** (2009) Effects of *Lactobacillus buchneri* and *Lactobacillus plantarum* on fermentation, aerobic stability, bacteria diversity and ruminal degradability of alfalfa silage. *World Journal of Microbiology and Biotechnology* **25**, 965–971.
- Zhang W, Grimi N, Jaffrin MY, Ding L and Tang B** (2017) A short review on the research progress in alfalfa leaf protein separation technology. *Journal of the Society of Chemical Industry* **92**, 2894–2900.
- Zielińska K, Fabiszewska A and Stefańska I** (2015) Different aspects of *Lactobacillus* inoculants on the improvement of quality and safety of alfalfa silage. *Chilean Journal of Agricultural Research* **75**, 298–306.